

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 May 2002 (02.05.2002)

PCT

(10) International Publication Number
WO 02/34722 A1

(51) International Patent Classification⁷: C07D 233/54,
C07F 5/02, C07C 307/02, C08F 4/00, 36/04

(74) Agent: JACOBSON, Claude; Cabinet Lavoix, 2, place
d'Estienne-d'Orves, F-75441 Paris Cedex 09 (FR).

(21) International Application Number: PCT/IB01/02003

(22) International Filing Date: 25 October 2001 (25.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
00403008.6 27 October 2000 (27.10.2000) EP

(71) Applicant (for all designated States except US): CENTRE
NATIONAL DE LA RECHERCHE SCIENTIFIQUE
(C.N.R.S.) [FR/FR]; 3, rue Michel Ange, F-75016 Paris
(FR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,
ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(72) Inventors; and

(75) Inventors/Applicants (for US only): CHAUVIN, Yves
[FR/FR]; 10, Place François Sicard, F-37000 Tours (FR).
MAGNA, Lionel [FR/FR]; Le Fenouillet, Les Jonquilles
C1, F-83400 Hyeres (FR). NICCOLAI, Gerald, Peter
[FR/FR]; 1, rue du Professeur Calmette, F-69100 Villeur-
banne (FR). BASSET, Jean-Marie [FR/FR]; 18, Chemin
J.B. Gilliard, F-69300 Caluire (FR).

WO 02/34722 A1

(54) Title: IMIDAZOLIUM SALTS AND THEIR USE OF THESE IONIC LIQUIDS AS A SOLVENT

(57) Abstract: New 1,2,3- or 1,2,3,4- or 1,2,3,4,5- substituted imidazolium salts and their uses as solvent in catalysed organic reactions, as well as compositions containing them and a transition metal compound. They can be used in the following reactions: telomerisation of conjugated dienes, dimerisation of olefins, oligomerisation of olefins, polymerization of olefins, alkylation of olefins, hydrogenations of olefins, olefin metathesis, hydroformylation of olefins, ring-closing metathesis of olefins, ring-opening metathesis polymerisation of olefins, symmetric or asymmetric epoxidation of olefins (including heteroatom substituted olefins) and cyclopropanation of olefins, condensation reaction, hydrogenation reactions, isomerisation reaction, Suzuki cross-coupling reactions, amination reaction, partial oxidation of alkanes, kinetic resolution of racemic mixtures, hydrogenation of imines, hydrogenation of ketones, transfer hydrogenation and hydroxylation of aromatic organic compounds.

IMIDAZOLIUM SALTS AND THEIR USE OF THESE IONIC LIQUIDS AS A SOLVENT

5 [0001] The present invention relates to the use of imidazolium salts as solvents and to new imidazolium salts and compositions containing the same. In particular the imidazolium salts may be used as a solvent in organic reactions, in particular in catalytic reactions such as two-phase catalytic conversions of olefins.

[0002] Low melting salts, which are liquid at room temperature up to 200 °C, can
10 be used as liquid media in many organic reactions. Usually these salts are referred to as "ionic liquids". Frequently the ionic liquids have two functions in that they not only serve as a solvent for the reaction but also as a catalyst or catalyst component.

[0003] In other cases the ionic liquids serve only as a solvent for the catalyst, that is the ionic liquid is not miscible with the reactants or the reaction product. In this case
15 the reaction occurs at the phase boundary of the catalyst solution and the reaction products form a separate phase. This process using two separate liquid phases is suitable, if the reaction product must be quickly removed from the reaction mixture in order to avoid subsequent reactions. Moreover the catalyst and the reaction product can be separated under gentle conditions.

20 [0004] Further advantages of ionic liquids are their chemical and thermal stability, which allows their application in many processes. Further they are not volatile and therefore they are safer for chemical industry personnel and the general public and environment.

[0005] Among the ionic liquids known as solvents in the art imidazolium salts have
25 attracted attention (for a recent review, see as an example "Ionic liquids" by J.D. Holbrey and K. R. Seddon in *Clean Products and Processes*, 1, (1999), 223-236).

[0006] For example *CHEMTECH*, (September 1995), pages 26 and following pages discloses a mixture of 1,3-dialkylimidazolium chloride, in particular 1-n-butyl-3-methyl-imidazolium chloride (abbreviated as BMI⁺Cl⁻) and aluminium chloride
30 and/or ethyl aluminium chloride as non-aqueous solvent for the catalyst.

[0007] *Am. Chem. Soc., Div. Pet. Chem.*, (1992), 37, pages 370 *sqq.* discloses the dimerisation of propene in the presence of a solution of $\text{NiCl}_2(\text{PR}_3)_2$, ($\text{R}=\text{i-C}_3\text{H}_7$) in a mixture of BMI^+Cl^- and AlCl_3 as ionic liquid.

[0008] EP-A-0 776 880 (corresponding to US 5,874,638) discloses for example a process for the hydroformylation of olefinic compounds in the presence of an ionic liquid.

[0009] *Angew. Chem.*, (1995), 107(23/24), pages 2941 *sqq.* also discloses the hydroformylation reaction using liquid 1,3-dialkylimidazolium salts.

[0010] British patent application GB-A-2 150 740, as well as EP-A-0 404 179 and EP-A-0 398 358 disclose 1,2,3-trialkylimidazolium halides and their mixtures with aluminium halides as electrolytes. WO 99/40025 also discloses quaternary onium salts useful as electrolytic solutes. Still other heterocyclic onium salts may be for example used in non-aqueous batteries as disclosed in US 5,827,602 and US 5,683,832.

[0011] Other uses of quaternary imidazolium salts are also known, for example as microbicides or plant growth regulators in FR-A-2 434 156, FR-A-2 380 732, FR-A-2 302 301 and FR-A-2 303 802, and as anti-static agents for synthetic fibres in JP 06 812354.


[0012] Haloiminium salts (e.g. tri-substituted imidazolium chloride) have been reported as being useful for lactic acid polymerisation (see JP 01 135662). WO 95/21871 presents disubstituted imidazolium halides as catalysts in olefinic polymerisation reactions.

[0013] German patent application no. 199 19 494.7 (corresponding to international patent application WO 00/66597) discloses non aqueous ionic ligand liquids of the formula $(\text{Q}_1^+)_b\text{A}'^c$, wherein Q_1^+ is a singly charged ammonium cation, optionally substituted with organic group(s), or the equivalent of a multiply charged ammonium cation and A'^c is an anion of a sulphonated or carboxylated triester of phosphorous acid and c is an integer of at least one. Among the ammonium cations there are also mentioned 1,2,3,4,5-pentamethylimidazolium, 1,2,3,5-tetramethyl-4*H*-imidazolium, 1,2,3,4-tetramethyl-5*H*-imidazolium, and 1-trimethylsilyl-2,3,5-trimethyl-4*H*-imidazolium.

- [0014] J. Dupont et al. (*Organometallics*, (1998), 17, 815-819) disclose a catalytic process of hydrodimerisation of 1,3-butadiene using palladium compounds dissolved in ionic liquids. As palladium catalyst compounds $[(\eta^3\text{-C}_4\text{H}_7)\text{Pd-}\mu\text{-Cl}]_2$, $[(\eta^3\text{-C}_4\text{H}_7)\text{Pd}(1,5\text{-cyclooctadiene})]$ $[\text{BF}_4]$ and palladium acetate have been used which are told not to be completely soluble and stable in the ionic liquids 1-n-butylmethylimidazolium tetrafluoroborate ($\text{BMI}^+\text{.BF}_4^-$) and 1-n-butylmethylimidazolium hexafluorophosphate ($\text{BMI}^+\text{.PF}_6^-$) at room temperature. At the end of the performed hydrodimerisation reactions however, metallic palladium was detected thus limiting the reutilisation of the catalytic system. This has been attributed to the instability of these catalysts to water. The formation of metallic palladium could be suppressed by the use of a new catalyst precursor $(\text{BMI})_2\text{PdCl}_4$ which has been obtained by reacting PdCl_2 with a 2 molar excess of 1-n-butyl-3-methylimidazolium chloride in acetonitrile at reflux temperature. However even for this stable catalyst, conversions reported were low.
- [0015] In an attempt to use conventional palladium phosphine catalysts in the presence of ionic liquids such as the above mentioned 1-n-butyl-methylimidazolium tetrafluoroborate ($\text{BMI}^+\text{.BF}_4^-$) or 1-ethyl-3-methylimidazolium bis(trifluoromethanesulphonyl)imide ($\text{EMI}^+\text{.Tf}_2\text{N}^-$) for the telomerisation of butadiene for example with methanol, the present inventors found surprisingly that the conventional palladium phosphine catalyst system in the presence of ionic 1,3-dialkylimidazolium liquids in contrast to the above mentioned catalyst compounds like $[(\eta^3\text{-C}_4\text{H}_7)\text{Pd-}\mu\text{-Cl}]_2$, $[(\eta^3\text{-C}_4\text{H}_7)\text{Pd}(1,5\text{-cyclooctadiene})]$ $[\text{BF}_4]$ or palladium acetate shows almost no reactivity, and thus was not available for a telomerisation process in the presence of the ionic liquid.
- [0016] Therefore, the object underlying the present invention was to find new stable imidazolium salts which can be used, in particular, in catalytic processes without having a detrimental effect on the catalytic process. Surprisingly the applicant found that imidazolium salts wherein at least the 1,2 and 3-position of the imidazolium heterocycle are substituted have a less detrimental effect when used as a solvent and therefore could be advantageously used as a solvent in catalysed reactions, for example in the catalytic telomerisation of conjugated dienes.

[0017] Thus in accordance with the present invention there is provided a new use of 1,2,3-substituted imidazolium salts, 1,2,3,4-substituted imidazolium salts or 1,2,3,4,5-substituted imidazolium salts, with the exception of 1,2,3,4,5-pentamethylimidazolium-, 1,2,3,5-tetramethyl-4*H*-imidazolium-, 1,2,3,4-tetramethyl-5*H*-imidazolium-, and 1-trimethylsilyl-2,3,5-trimethyl-4*H*-imidazolium-salts with
 5 sulphonated or carboxylated triesters of phosphorous acid as anion, as solvent.

[0018] Preferred 1,2,3-substituted imidazolium salts, 1,2,3,4-substituted imidazolium salts or 1,2,3,4,5-substituted imidazolium salts are those of the following formula (1):



(1)

10

wherein R¹ and R³ are the same or different and are each selected from the group consisting of a chiral or achiral alkyl group having up to 10, preferably up to 6 carbon atoms, a chiral or achiral cycloalkyl group of 4 to 10 carbon atoms, a chiral or
 15 achiral aryl group of 6 to 10 carbon atoms, and a chiral or achiral tri(C1-C10)alkylsilyl group,

R² is selected from the group consisting of a chiral or achiral alkyl group having up to 10, preferably up to 6 carbon atoms, a chiral or achiral cycloalkyl group of 4 to 10 carbon atoms, a chiral or achiral aryl group of 6 to 10 carbon atoms and a
 20 halogen atom; more preferably R² is an alkyl group, a cycloalkyl group or an aryl group as defined above,

R⁴ and R⁵ are the same or different and are each selected from the group consisting of hydrogen, a chiral or achiral alkyl group having up to 10, preferably up to 6 carbon atoms, a chiral or achiral cycloalkyl group of 4 to 10 carbon atoms, a
 25 chiral or achiral aryl group of 6 to 10 carbon atoms, a chiral or achiral tri(C1-C10)alkylsilyl group, and a halogen atom; more preferably R⁴ and R⁵ are hydrogen, an alkyl group, a cycloalkyl group, an aryl group, or a

tri(C1-C10)alkylsilyl group, still more preferably hydrogen,

and X⁻ represents an anion.

[0019] By "chiral or achiral" is meant groups that may eventually possess one or more chiral centres resulting in an optical activity and/or groups that have an intrinsic optical activity.

[0020] The aforementioned alkyl group having up to 10 carbon atoms (sometimes abbreviated as "C1-C10"), preferably up to 6 carbon atoms (sometimes abbreviated as "C1-C6") in the definitions of R¹ to R⁵ includes optionally substituted linear or branched alkyl groups, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, *iso*-butyl, *sec*-butyl, *tert*-butyl, n-pentyl, *iso*-pentyl, *neo*-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl and n-decyl. Alkyl groups may be substituted by one or more substituents, identical or different. Among these, preferred substituents are halogen atoms. Preferred are alkyl groups with up to 6 carbon atoms, e.g. methyl, ethyl, propyl, *iso*-propyl, butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, 1-pentyl, 2-pentyl, 3-pentyl, *iso*-amyl, n-hexyl and *iso*-hexyl. Preferred substituted alkyl are haloalkyl and perhaloalkyl, e.g. trifluoromethyl, 1,1,1-trifluoroethyl, perfluoroethyl, perfluoropropyl, 1,1,1,2,2,3,3-heptafluorobutyl and perfluorobutyl.

[0021] The aforementioned cycloalkyl group of 6 to 10 carbon atoms includes for example cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclo-octyl, as well as bicycloalkyl groups, e.g. bicyclo[3.3.0]octyl, bicyclo[4.3.0]nonyl, bicyclo[4.4.0]decyl and bicyclo[2.2.1]heptyl. Tricycloalkyl groups, for example adamantyl, are included in the definition of cycloalkyl and are also part of the present invention. Cycloalkyl groups may be substituted by one or more identical or different substituents chosen, for example, from among alkyl groups and halogen atoms.

[0022] The aforementioned aryl group of 6 to 10 carbon atoms includes for example phenyl or naphthyl. Aryl groups may bear one or more substituents, identical or different.

[0023] The skilled artisan will appreciate that all various substituents on the alkyl, cycloalkyl and/or aryl groups are all identical or different and, when present, serve to the formation of tri-, tetra-, and penta-substituted imidazolium salts, the use of which being one of the purposes of the present invention.

[0024] With respect to the above mentioned (C1-C6)alkyl moiety of the tri(C1-C6)alkylsilyl group it is referred to the definition of the (C1-C6)alkyl groups for R¹ to R⁵.

[0025] Halogen atoms in the above definitions include for example fluorine, chlorine, bromine and iodide.

[0026] In the above formula (1), X⁻ anion represents any anion known by the skilled artisan.

[0027] Preferred anions X⁻ in the above mentioned formula (1) are preferably selected from, but not limited to, anions of the group consisting of fluoride, chloride, bromide, iodide, hexafluorophosphate, hexafluoroantimonate, hexafluoroarsenate, fluorosulphonate, tetrafluoroborate, bis-perfluoro(C1-C10)alkylsulphonyl amides, perfluoro(C1-C10)alkyl sulphonates, dichlorocuprate, tetrachloroborate, tetrachloroaluminate, heptachlorodialuminate (Al₂Cl₇⁻), trichlorozincate, and anions of sulphonated or carboxylated (alkyl, cycloalkyl and/or aryl)-phosphines or -phosphites.

[0028] Particularly preferred anions in the above formula (1) are those selected from fluoride, chloride, bromide, iodide, hexafluorophosphate, fluorosulphonate, tetrachloroborate, tetrafluoroborate, bis-perfluoro(C1-C10)alkylsulphonyl amides, perfluoro(C1-C10)alkyl sulphonates, and anions of sulphonated or carboxylated (alkyl, cycloalkyl and/or aryl)-phosphines or -phosphites.

[0029] With respect to the aforementioned (C1-C10)alkyl moiety in the definitions "bis-perfluoro(C1-C10)alkylsulphonyl amide" and "perfluoro(C1-C10)alkyl sulphonate" it is referred to the definition of the C1-C10 alkyl groups for R¹ to R⁵ above.

[0030] Preferred imidazolium salts of formula (1) used as solvents are those having the following characteristics taken alone or in combination:

- R¹, R² and R³, identical or different, are each independently chosen from (C1-C6)alkyl, (C1-C6)haloalkyl and cycloalkyl;
- R⁴ and R⁵, identical or different, are each independently chosen from hydrogen, (C1-C6)alkyl, (C1-C6)haloalkyl and cycloalkyl;
- and X⁻ represents an anion as hereinbefore described.

[0031] Most preferred imidazolium salts of formula (1) used as solvents have the following characteristics taken alone or in combination:

- R^1 , R^2 and R^3 , identical or different, are each independently chosen from methyl, ethyl, propyl, *iso*-propyl, butyl, *sec*-butyl, 1-pentyl, 2-pentyl, 3-pentyl, *iso*-pentyl, hexyl, *iso*-hexyl, cyclohexyl, trifluoromethyl, 1,1,1-trifluoroethyl, perfluoroethyl, perfluoropropyl, 1,1,1,2,2,3,3-heptafluorobutyl, and perfluorobutyl;
- R^4 and R^5 , identical or different, are each independently chosen from hydrogen, methyl, ethyl, propyl, *iso*-propyl, butyl, *sec*-butyl, 1-pentyl, 2-pentyl, 3-pentyl, *iso*-pentyl, hexyl, *iso*-hexyl, cyclohexyl, trifluoromethyl, 1,1,1-trifluoroethyl, perfluoroethyl, perfluoropropyl, 1,1,1,2,2,3,3-heptafluorobutyl, and perfluorobutyl;
- and X^- represents an anion as hereinbefore described.

[0032] Particular preferred solvents are 1,2,3-tri(C1-C10)alkyl-imidazolium salts. With respect to the aforementioned (C1-C10)alkyl moiety in the "1,2,3-tri(C1-C10)-alkyl-imidazolium salts" it is referred to the definition of the C1-C10 alkyl groups for R^1 to R^5 above.

[0033] (Alkyl, cycloalkyl and/or aryl)-phosphines or phosphites in the definition of the anions of sulphonated or carboxylated (alkyl, cycloalkyl and/or aryl)-phosphines or phosphites include all kinds of anions of alkyl and aryl phosphines or phosphites like monodentate or bidentate phosphines or phosphites having up to 36 carbon atoms in the hydrocarbon moieties, like alkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, etc.

[0034] Especially preferred salts of formula (1) used as solvents include 1-ethyl-2,3-dimethylimidazolium bromide ([EMMI][Br]), 1-ethyl-2,3-dimethylimidazolium bistrifluoromethylsulphonylamide ([EMMI][Tf₂N]), 1-ethyl-2,3-dimethylimidazolium trifluoromethylsulphonate ([EMMI][CF₃SO₃]), 1-n-butyl-2,3-dimethylimidazolium chloride ([BMMI][Cl]), 1-n-butyl-2,3-dimethylimidazolium hexafluorophosphate ([BMMI][PF₆]), 1-n-butyl-2,3-dimethylimidazolium bistrifluoromethylsulphonylamide ([BMMI][Tf₂N]), and 1-n-butyl-2,3-methylimidazolium tetrafluoroborate ([BMMI][BF₄]).

[0035] In a particular preferred embodiment, 1-n-butyl-2,3-dimethylimidazolium salt is used as a solvent, and more specifically 1-n-butyl-2,3-dimethylimidazolium

chloride, 1-n-butyl-2,3-dimethylimidazolium hexafluorophosphate, 1-n-butyl-2,3-dimethylimidazolium bistrifluoromethylsulphonylamide and/or 1-n-butyl-2,3-methylimidazolium tetrafluoroborate are used as solvents.

[0036] Preferably the above substituted imidazolium salts are used as a solvent in organic reactions, in particular in the presence of a catalyst, wherein the catalyst may
5 comprise a transition metal compound, in particular from the group VIIB of the periodic system of elements.

[0037] Preferably the above substituted imidazolium salts are used as a solvent in organic reactions, selected from the group consisting of the catalytic conversion of
10 olefins selected from the telomerisation of conjugated dienes, the dimerisation of olefins, the oligomerisation of olefins, the polymerisation of olefins, the alkylation of olefins, the hydrogenation of olefins, the olefin metathesis, the hydroformylation of olefins, the ring-closing metathesis of olefins, the ring-opening metathesis polymerisation of olefins, the symmetric or asymmetric epoxidation of olefins
15 (including heteroatom substituted olefins) and the cyclopropanation of olefins, the condensation reaction, the hydrogenation of, in particular, aldehydes, the isomerisation reaction, the Suzuki cross-coupling reactions, the amination reaction, the partial oxidation of alkanes, the kinetic resolution of racemic mixtures, the hydrogenation of imines, the hydrogenation of ketones, the transfer hydrogenation,
20 and the hydroxylation of aromatic organic compounds.


[0038] Preferred reactions, wherein the solvent is an imidazolium salt of formula (1), are reactions of catalytic conversion of olefins.

[0039] A particular preferred process is the telomerisation of conjugated dienes.

[0040] In some cases the 1,2,3-substituted imidazolium salts, 1,2,3,4-substituted
25 imidazolium salts or 1,2,3,4,5-substituted imidazolium salts, or mixtures thereof, with the exception of 1,2,3,4,5-pentamethyl imidazolium-, 1,2,3,5-tetramethyl-4H-imidazolium-, 1,2,3,4-tetramethyl-5H-imidazolium-, and 1-trimethylsilyl-2,3,5-trimethyl-4H-imidazolium-salts with sulphonated or carboxylated triesters of phosphorous acid as anion, act also as a catalyst, and accordingly, the present
30 invention is also related to the use of the above imidazolium salts as a catalyst or catalyst component, in particular in transition metal catalyst systems. This is

particularly true in those cases where the anion of the imidazolium salt is a phosphorous containing anion like anions of the aforementioned sulphonated or carboxylated (alkyl and/or aryl) phosphines or phosphites.


[0041] Such anions are disclosed for example in EP-A-0 924 218 (US 6,103,908) and shall be included within the scope of the present invention. EP-A-0 924 218 discloses nonaqueous ionic ligand liquids of the formula $(Q^+)_a A^{a-}$, wherein Q^+ is a singly charged quaternary ammonium and/or phosphonium cation or the equivalent of a multiply charged ammonium and/or phosphonium cation and A^{a-} is a triarylphosphine anion of the formula :



where Ar_1 , Ar_2 and Ar_3 are individually aryl of 6 to 14 carbon atoms, the substituents Y_1 , Y_2 and Y_3 are individually selected from the group consisting of alkyl and alkoxy of 1 to 4 carbon atoms, chlorine, bromine, hydroxyl, cyano, nitro and amino groups of the formula NR^6R^7 , where R^6 and R^7 are individually hydrogen or alkyl of 1 to 4 carbon atoms, m_1 , m_2 and m_3 are individually integers from 0 to 5, n_1 , n_2 and n_3 are individually integers from 0 to 3, where at least one of n_1 , n_2 and n_3 is equal to or greater than 1, and a is $n_1+n_2+n_3$, and amines and/or phosphines derived from Q^+ are present in an excess of up to 5 equivalents over the stoichiometrically required amount for the formation of $(Q^+)A^{a-}$ or alkali metal or alkaline earth metal salts of the triarylphosphines A^{a-} are present in an excess of up to 5 equivalents over the stoichiometrically required amount for the formation of $(Q^+)A^{a-}$.

[0042] Similarly the above mentioned German patent application no. 19919494.7 (corresponding to international patent application no. WO 00/66597) discloses

suitable non-aqueous ionic ligand liquids of the formula $(Q_1^+)_b A'^b$, wherein Q_1^+ is a singly charged ammonium cation, optionally substituted with organic group(s), or the equivalent of a multiply charged ammonium cation and A'^b is an anion of a sulphonated or carboxylated triester of phosphorous acid and b is an integer of at least one. These ammonium salts of sulphonated or carboxylated phosphorous acids may be formally derived from phosphorous acid by esterification with the ammonium salts of hydroxysulphonic acids or hydroxycarboxylic acids of the following general formula



wherein ac is the acid residue, namely the sulphonic acid residue $-SO_3^-$, and the carboxylic acid residue, respectively and Q is as already mentioned a singly charged ammonium cation, optionally substituted with organic group(s), or the equivalent of a multiply charged ammonium cation, Y represents an organic residue and is preferably a branched or non-branched saturated aliphatic hydrocarbon group having a total of up to 20 carbon atoms, optionally substituted by hydroxy or C1-C10 alkoxy, c and d are integers of at least 1, d is preferably 1 or 2. Also these anions of sulphonated or carboxylated phosphorous acids, in particular sulphonated aryl or carboxylated aryl phosphorous acids may be used as anions of the imidazolium cations in accordance with the present invention.


[0043] Further the present invention relates to the use of the above mentioned 1,2,3-substituted imidazolium salts, 1,2,3,4-substituted imidazolium salts or 1,2,3,4,5-substituted imidazolium salts, or mixtures thereof, with the exception of 1,2,3,4,5-pentamethylimidazolium-, 1,2,3,5-tetramethyl-4*H*-imidazolium-, 1,2,3,4-tetramethyl-5*H*-imidazolium-, and 1-trimethylsilyl-2,3,5-trimethyl-4*H*-imidazolium-salts with a sulphonated or carboxylated triester of phosphorous acid as anion, in a two-phase catalytic reaction, preferably as solvent and/or catalyst or catalyst component.

[0044] The present invention also relates to the use of 1,2,3- and/or 1,2,3,4- and/or 1,2,3,4,5-substituted imidazolium salts, in particular those defined above, with a metal compound catalyst or catalyst component, in particular in organic reactions as

defined above. The salts can have a role of solvent and possibly of catalyst or catalyst component, preferably solvent.

[0045] The use of salts containing chiral group(s) leading to a chiral solvent may advantageously confer enantioselectivity to the reaction wherein they are used.

5 [0046] Still further the present invention relates to new imidazolium salts of the following formula (1):



10 wherein R¹ and R³ are the same or different and are each selected from the group consisting of a chiral or achiral alkyl group having up to 10, preferably up to 6 carbon atoms, a chiral or achiral cycloalkyl group of 4 to 10 carbon atoms, a chiral or achiral aryl group of 6 to 10 carbon atoms, and a chiral or achiral tri(C1-C10)alkylsilyl group,

15 R² is selected from the group consisting of a chiral or achiral alkyl group having up to 10, preferably up to 6 carbon atoms, a chiral or achiral cycloalkyl group of 4 to 10 carbon atoms, a chiral or achiral aryl group of 6 to 10 carbon atoms and a halogen atom; more preferably R² is an alkyl group, a cycloalkyl group or an aryl group as defined above,

20 R⁴ and R⁵ are the same or different and are each selected from the group consisting of hydrogen, a chiral or achiral alkyl group having up to 10, preferably up to 6 carbon atoms, a chiral or achiral cycloalkyl group of 4 to 10 carbon atoms, a chiral or achiral aryl group of 6 to 10 carbon atoms, a chiral or achiral tri(C1-C10)alkylsilyl group, and a halogen atom; more preferably R⁴ and R⁵ are
25 hydrogen, an alkyl group, a cycloalkyl group, an aryl group, or a tri(C1-C10)alkylsilyl group, still more preferably hydrogen,

and X⁻ represents an anion,

with the exception of 1,2,3,4,5-pentamethylimidazolium-, 1,2,3,5-tetramethyl-4*H*-imidazolium-, 1,2,3,4-tetramethyl-5*H*-imidazolium, and 1-trimethylsilyl-2,3,5-trimethyl-4*H*-imidazolium-salts of sulphonated or carboxylated triesters of phosphorous acid, 1,2,3,4,5-pentamethylimidazolium iodide, 1,2,3-trimethylimidazolium bromide, 1,2,3-trimethylimidazolium iodide, 1,2-dimethyl-3-ethylimidazolium bromide, 1,2-dimethyl-3-ethylimidazolium chloride, 1,2-dimethyl-3-butylimidazolium fluoride, 1-butyl-2,3-dimethylimidazolium salt of 2,2'-(2,5-cyclohexadiene-1,4-ylidene)bis[propanedinitrile], 1,3-dimethyl-2,4,5-trichloroimidazolium tetrafluoroborate, 1,3-dimethyl-2,4,5-tribromoimidazolium tetrafluoroborate, 1,3-dimethyl-2,4,5-tribromoimidazolium bromide.

[0047] With respect to the explanation of the substituents it is referred to the definitions and preferred definitions given above.

[0048] Preferred imidazolium salts according to the invention are those having the following characteristics taken alone or in combination:

- R^1 , R^2 and R^3 , identical or different, are each independently chosen from (C1-C6)alkyl, (C1-C6)haloalkyl and cycloalkyl;
- R^4 and R^5 , identical or different, are each independently chosen from hydrogen, (C1-C6)alkyl, (C1-C6)haloalkyl and cycloalkyl;
- and X^- represents an anion as hereinbefore described.

[0049] Most preferred imidazolium salts of formula (1) have the following characteristics taken alone or in combination:

- R^1 , R^2 and R^3 , identical or different, are each independently chosen from methyl, ethyl, propyl, *iso*-propyl, butyl, *sec*-butyl, 1-pentyl, 2-pentyl, 3-pentyl, *iso*-pentyl, hexyl, *iso*-hexyl, cyclohexyl, trifluoromethyl, 1,1,1-trifluoroethyl, perfluoroethyl, perfluoropropyl, 1,1,1,2,2,3,3-heptafluorobutyl, and perfluorobutyl;
- R^4 and R^5 , identical or different, are each independently chosen from hydrogen, methyl, ethyl, propyl, *iso*-propyl, butyl, *sec*-butyl, 1-pentyl, 2-pentyl, 3-pentyl, *iso*-pentyl, hexyl, *iso*-hexyl, cyclohexyl, trifluoromethyl, 1,1,1-trifluoroethyl, perfluoroethyl, perfluoropropyl, 1,1,1,2,2,3,3-heptafluorobutyl, and perfluorobutyl;
- and X^- represents an anion as hereinbefore described.

[0050] Particular preferred salts of the present invention are 1,2,3-tri(C1-C10)alkyl-imidazolium salts. With respect to the aforementioned (C1-C10)alkyl moiety in the "1,2,3-tri(C1-C10)alkyl-imidazolium salts" it is referred to the definition of the C1-C10 alkyl groups for R¹ to R⁵ above.

5 [0051] (Alkyl, cycloalkyl and/or aryl)-phosphines or phosphites in the definition of the anions of sulphonated or carboxylated (alkyl, cycloalkyl and/or aryl)-phosphines or phosphites include all kinds of anions of alkyl and aryl phosphines or phosphites like monodentate or bidentate phosphines or phosphites having up to 36 carbon atoms in the hydrocarbon moieties, like alkyl, aryl, arylalkyl, cycloalkyl,
10 cycloalkylalkyl, etc.

[0052] In a preferred embodiment, salts of formula (1) include those wherein R¹, R², R³, R⁴ and R⁵ are as described above and X⁻ is chosen from fluoride (F⁻), chloride (Cl⁻), bromide (Br⁻), iodide (I⁻), bistrifluoromethylsulphonylamide (Tf₂N⁻), trifluoromethylsulphonate (CF₃SO₃⁻), hexafluorophosphate (PF₆⁻), and
15 tetrafluoroborate (BF₄⁻).

[0053] More specifically, X⁻ is chosen from bistrifluoromethylsulphonylamide (Tf₂N⁻), trifluoromethylsulphonate (CF₃SO₃⁻), hexafluorophosphate (PF₆⁻), and tetrafluoroborate (BF₄⁻).

[0054] Especially preferred salts of formula (1) used as solvents include 1-ethyl-
20 2,3-dimethylimidazolium bromide ([EMMI][Br]), 1-ethyl-2,3-dimethylimidazolium bistrifluoromethylsulphonylamide ([EMMI][Tf₂N]), 1-ethyl-2,3-dimethylimidazolium trifluoromethylsulphonate ([EMMI][CF₃SO₃]), 1-n-butyl-2,3-dimethylimidazolium chloride ([BMMI][Cl]), 1-n-butyl-2,3-dimethylimidazolium hexafluorophosphate ([BMMI][PF₆]), 1-n-butyl-2,3-dimethylimidazolium
25 bistrifluoromethylsulphonylamide ([BMMI][Tf₂N]), and 1-n-butyl-2,3-dimethylimidazolium tetrafluoroborate ([BMMI][BF₄]).

[0055] Particular preferred imidazolium salts of the invention are 1-n-butyl-2,3-dimethylimidazolium bistrifluoro(C1-C10, preferably C1 (methyl))alkylsulphonylamide, 1-n-butyl-2,3-dimethylimidazolium tetrafluoroborate and 1-n-butyl-2,3-
30 dimethylimidazolium hexafluorophosphate.

[0056] The present invention also concerns compositions or mixtures comprising at least one 1,2,3- or 1,2,3,4- or 1,2,3,4,5-substituted imidazolium salt, in particular as defined above, as a solvent; and at least one transition metal or at least one catalyst, in particular a catalyst comprising a transition metal compound; transition metal may be in particular from group VIIIB, e.g. as further defined herein. The catalyst can be any catalyst usually used in the catalytic reactions defined above. The composition may also comprise the reactant(s).

[0057] These compositions or mixtures may comprise salts as defined above with the specified both kind of exceptions.

[0058] Such compositions or mixtures may also comprise a phosphorous-containing compound as defined hereinafter.

[0059] The amount of these phosphorus-containing compounds is in general from 0.01 to 100, preferably from 1 to 20, in particular from 1 to 5 mole per mole of the transition metal.

[0060] The concentration of the transition metal is usually in the range of 1 to 200 mM per litre of ionic liquid and more preferably 10 to 50.


[0061] The amount of the salt in the reaction system is usually 5 to 100 parts per 100 parts of the reactant(s), e.g. diolefin, preferably 10 to 50.

[0062] Usually the imidazolium salts have a melting point of below 200 °C, preferably below 160 °C, more preferably below 140°C, still more preferably below 120°C, and most preferably below 100 °C, in particular, below 90 °C (the melting points are measured at normal pressure with a digital apparatus such as of Electrothermal).

[0063] 1-n-Butyl-2,3-dimethylimidazolium chloride (BMMI^+Cl^-) has a melting point of 104-105 °C, 1-n butyl-2,3-dimethylimidazolium bistrifluoromethylsulphonylamide ($\text{BMMI}^+\text{Tf}_2\text{N}^-$), and 1-n-butyl-2,3-dimethylimidazolium tetrafluoroborate ($\text{BMMI}^+\text{BF}_4^-$) are liquids at room temperature.

[0064] Synthesis of non aqueous ionic liquids based on unsymmetrical substituted 1,2,3-trialkylimidazolium can be performed following two main schemes. According to the first one, the N-alkylimidazole or 1,2-dialkylimidazole (generally commercial products) are alkylated with an organic halide and the desired salt is then obtained by

anion exchange reactions (metathesis). According to the second one, the liquid is obtained directly by the quaternisation of an N-alkylimidazole, with the alkylating agent itself providing a suitable anion.



5

[0065] For the indirect route (1) one can refer to P. Bonhôte, A.-P. Dias, N. Papageorgiou, K. Kalyanasundaram, Gratzel, *M. Inorg. Chem.*, (1996), **35**, 1168-1178; J. S. Wilkes, M. J. Zaworotko, *J. Chem. Soc., Chem. Commun.*, (1992), 965-967; J. Fuller, R. T. Carlin, R. A. Osteryoung, *J. Electrochem. Soc.*, (1997), **144**, 3881-3886. Generally, bromide or chloride imidazolium can be used as intermediate in the synthesis of the final ionic liquid. The anion exchange involved in the second step can be performed in different solvents: water (see J. D. Holbrey, K. R. Seddon, *J. Chem. Soc., Dalton Trans.*, (1999), 2133-2139) or acetone (see P. A. Suarez, J. E. L. Dullius, S. Einloft, R. F. de Souza, J. Dupont, *Polyhedron*, (1996), **15**, 1217-1219 and P. A. Suarez, S. Einloft, J. E. L. Dullius, R. F. de Souza, J. Dupont, *J. Chim. Phys.*, (1998), **95**, 1626-1639).

[0066] The direct synthesis route (2) allows one to obtain an ionic liquid without any addition of halide by the reaction of an N-alkylimidazole with an appropriate alkylating agent. The alkylating agent must be able to realise both the quaternisation of the imidazole ring and the introduction of a non-coordinating anion. Can be used among others, alkylating agents derived from alkyltriflates (methyl and ethyltriflate) (see P. Bonhôte, A.-P. Dias, N. Papageorgiou, K. Kalyanasundaram, Gratzel, *M. Inorg. Chem.*, (1996), **35**, 1168-1178). Reaction is carried out in refluxing 1,1,1-trichloroethane, a solvent chosen for its stability toward strongly alkylating agents

25

and the insolubility of the imidazolium salts in this milieu. To prevent alkyltriflate hydrolysis the reaction must be conducted under dry argon. Nearly quantitative yields were obtained.

[0067] The syntheses of $[R^1R^2I][BF_4]$ and $[R^1R^2I][PF_6]$ salts can be prepared using the first route which involves metathesis reaction from the corresponding chloride or bromide salts with $NaBF_4$ and $NaPF_6$ in water (see J. D. Holbrey, K. R. Seddon, *J. Chem. Soc., Dalton Trans.*, (1999), 2133-2139), methanol (see J. S. Wilkes, M. J. Zaworotko, *J. Chem. Soc., Chem. Commun.*, (1992), 965-967), or acetone (see P. A. Suarez, J. E. L. Dullius, S. Einloft, R. F. de Souza, J. Dupont, *Polyhedron*, (1996), 15, 1217-1219). They can also be obtained by using other alkylating agents. Those are triethyloxonium tetrafluoroborate and triethyloxonium hexafluorophosphate. As with ethyltriflate, the oxonium salt $[Et_3O][BF_4]$ reacts with one equivalent of 1-alkylimidazole (or 1,2-dialkylimidazole) in refluxing methylene dichloride with very high yields (see patent application by H. Olivier, F. Favre., IFP, no. FR 2.779.143, (1998)). All prepared imidazolium ionic liquids were air stable under ambient conditions and may be handled under normal laboratory conditions.

[0068] The use of the present invention is now exemplified further in a process for telomerising conjugated dienes.

[0069] The catalytic dimerisation of dienes under the concomitant addition of a nucleophilic reagent was reported simultaneously in 1967 by Smutny at Shell and Takahashi at Osaka University. The reaction is defined under the general term of telomerisation as the dimerisation of two conjugate diolefins (telomers) together with the addition of a third molecule (telogen) over one double bond equivalent.

[0070] Usually the process for telomerising a conjugated diene is carried out in the presence of a transition metal compound and a phosphorus-containing compound.

[0071] The conjugated diene preferably has 4 to 6 carbon atoms. Particularly preferred the conjugated diene is selected from butadiene, isopropene or 1,3-pentadiene. The most preferred conjugated diene is butadiene.

[0072] Telogens can be for example compounds containing active hydrogen, preferably a compound of the general formula

H-A

wherein A is selected from the group consisting of hydroxy, alkoxy, cycloalkoxy, hydroalkoxy, alkenyloxy, aryloxy, alkanoloxo, mono- or dialkylamino, tri(alkyl- and/or aryl)silyl, and an alkyl group substituted in the α -position with at least one electron attracting group, preferably selected from alkoxycarbonyl, alkanoyl and/or cyano, wherein the above mentioned alkyl moieties independently from each other are branched or linear and each may have up to 8 carbon atoms, and wherein the above mentioned aryl moieties independently from each other may have up to 10 carbon atoms.

[0073] Thus the compound containing active hydrogen includes for example water, alkanols like methanol, ethanol, n-propanol, *iso*-propanol, n-butanol, *iso*-butanol, *sec*-butanol etc., cycloalkanols like cyclohexanol etc., alkandiol like glycol, triethyleneglycol, polyethylene glycol, 1,2- or 1,3-propandiol etc., alkenols like allyl alcohol etc., aromatic alcohols like phenols, naphthols etc., carboxylic acids like acetic acid etc., amines like primary or secondary amines like methyl amine, ethyl amine, dimethyl amine, diethyl amine etc., tri(alkyl- and/or aryl) silanes, like trimethyl silane, triethyl silane, triphenyl silane, phenyl dimethyl silane etc., and CH-acidic compounds with activated methylene groups (so-called "Michael-Donors") like malonic acid dialkyl esters, acetic acid alkyl esters etc.

[0074] Particularly preferred telogens are water, methanol and glycol, and the most preferred is methanol.

[0075] The transition metal compound can be any transition metal compound suitable for catalysis of the telomerisation reaction in the presence of the phosphorous compound and the salt. Preferably it is a compound of an element of the group VIII B (or 8, 9 and 10) of the periodic system of elements, that is a compound selected from iron, cobalt, nickel, ruthenium, rhodium, palladium, osmium, iridium and platinum. More preferred compounds are selected from compounds such as cobalt, rhodium, nickel, platinum and platinum. Concerning possible transition metal compounds for carrying out the telomerisation process, reference is made to A. Behr in "Aspect of Homogeneous Catalysis, A series of Advances"; Edited by R. Ugo; D.

Reidel Publishing Company; Vol. 5; pp. 3-73, (1984) - Tsuji, *J. Adv. Organomet. Chem.*, (1979), 17, 141 - W. Keim, "Transition Metals in Homogeneous Catalysis", Ed. G. N. Schrauzer, p 59, Marcel Dekker, New York, (1971) - R. Baker, *Chem. Rev.*, 73, 487, (1973) - P. N. Rylander, "Organic Chemistry", vol. 28, p 175, Academic Press, New York, (1973).

[0076] Most preferred are palladium compounds, for example, known palladium compounds used in telomerisation processes for example those described in US 5,043,487, US 4,356,333, US 4,142,060, EP-A-0 436 226, WO 98/08794, WO 96/30636 and US 4,417,079 the relevant content of which is herein incorporated by reference.

[0077] Examples of these palladium compounds include soluble palladium(0) and palladium(II) compounds, for example, palladium acetylacetonate, π -allylpalladium acetate, π -allylpalladium chloride, palladium acetate, palladium carbonate, palladium nitrate, palladium chloride, sodium chloropalladate, bis(benzonitrile)palladium chloride, bis(triphenylphosphine)palladium chloride, bis(triphenylphosphine)-palladium acetate, bis(1,5-cyclooctadiene)palladium and bis(π -allyl)palladium.

[0078] A particular preferred palladium compound is palladium acetate ($\text{Pd}(\text{OAc})_2$).

[0079] The active species of the transition metal compound is a low-valence transition metal complex which may be formed by reduction in the presence of butadiene or by suitable reducing agents added.

[0080] The transition metal compound can be present in the reaction system in any optional amount but, from the standpoint of commercial production, the transition metal compound is preferably present in such an amount as to assure the concentration of transition metal atom of 0.1 to 50 mg atoms, more preferably 0.5 to 5 mg atoms per litre of the reaction mixture.

[0081] The ratio of the transition metal compound to the conjugated diene is not critical, but is preferably from 10^{-5} to 10^{-1} , in particular from 10^{-4} to 10^{-2} mole of the transition metal per mole of the conjugated diene.

[0082] The phosphorus-containing compound which forms part of the catalyst system is not particularly restricted but can be any phosphorus-containing compound, capable of coordinating to the transition metal compound, for example hydrophobic

or hydrophilic, water-soluble, mono- or bidentate phosphorus-containing compounds known for telomerisation processes (e.g. those known from US 5,043,487, US 5,345,007, US 4,356,333, EP-A-0 436 226, WO 98/08794, WO 95/30636, US 4,417,079 and US 4,142,060 the relevant content of which is herein incorporated by reference).

[0083] Suitable examples include phosphines or phosphites, preferably mono- or bidentate alkylphosphines, arylphosphines, arylalkylphosphines, alkylphosphites, arylphosphites, and arylalkylphosphites, wherein the hydrocarbon moieties may each independently have up to 36 carbon atoms, preferably 24 carbon atoms, more preferably 10 carbon atoms and may be substituted by one of three suitable substituents preferably selected from a sulphonic acid group or a salt thereof, a carboxylic acid group or a salt thereof and an alkyl group. Salts of the sulphonic acid or carboxylic acid group include for example alkali metal salts like sodium or potassium salts and ammonium salts.

[0084] Examples of these phosphines or phosphites include tributylphosphine, dimethyl-n-octylphosphine, tricyclohexylphosphine, triphenylphosphine (TPP), tritolylphosphine, tris(*para*-methoxyphenyl)phosphine, diphenylethylphosphine, dimethylphenylphosphine, 1,2-bis-diphenylphosphinoethane, triethylphosphite, tricyclohexylphosphite and triphenylphosphite, hydrophilic arylphosphines, like sulphonated or carboxylated arylphosphines, preferably water-soluble salts of mono-, di- or trisulphonated triphenylphosphine compounds like trisodium tris(*meta*-sulphonatophenyl)phosphine (TPPTS), bis(*para*-sulphonatophenyl)phenylphosphine dihydrate dipotassium salt (TPPDS) and diphenylphosphinobenzene-3-sulphonic acid sodium salt (TPPMS).

[0085] In general, the phosphines are preferred to the phosphites since the latter depending on the telogen used may undergo hydrolysis and rearrangement reactions. Particularly preferred are aryl phosphine compounds, sulphonated or not, like the above mentioned TPP, TPPTS, TPPDS and TPPMS.

[0086] Such phosphorous-containing compounds can be present in the compositions or mixtures defined above without being limited to a telomerisation process.

[0087] The amount of these phosphorus-containing compounds is in general from 1 to 100, preferably from 1 to 20, in particular from 1 to 5 mole per mole of the transition metal.

[0088] Depending on the telogen, to be telomerised, the phosphorus-containing compound used as the catalyst component and the relative amounts of the reactants in some instances monophasic reaction systems may be formed. These monophasic reaction systems can be usually transferred into biphasic systems by addition of at least one non-polar solvent (usually a solvent which is immiscible with water). Thereby a phase consisting of the solvent and the product (usually the upper phase) and a phase consisting of the salt (or ionic liquid) and the major part of catalyst (usually the lower phase) are formed. In some cases, depending on the telogen used and the catalyst system part of the catalyst may be contained also in the product phase. The non-polar solvent may include for example aliphatic hydrocarbons, for example alkanes like pentane, hexane, heptane, and octane etc., cycloalkanes like cyclopentane, cyclohexane etc., aromatic hydrocarbons like benzene, toluene, xylene etc. and aliphatic or aromatic ethers, like diethyl ether, tetrahydrofuran, anisol, methyl-*tert*-butyl ether (MTBE), diethylene glycol, dimethoxyethane, etc. The present invention includes also the case where the imidazolium salts are used together with such conventional non-polar solvents and/or polar solvents like water. A particular preferred non-polar solvent is n-heptane.

[0089] Depending on the telogen the desired products are for example *trans*- and *cis*-1-methoxy-2,7-octadiene and 3-methoxy-1,7-octadiene for methanol, *cis*- and *trans*-2,7-octadiene-1-ol and 1,7-octadiene-3-ol for water, and *cis*- and *trans*-1-hydroxy-2-(2,7-octadienyl-1-oxy)ethane and 1-hydroxy-2-(1,7-octadienyl-3-oxy)ethane for glycol.

[0090] The obtained unsaturated products can be hydrogenated catalytically into the corresponding saturated compounds in a known manner.

[0091] The telomerisation reaction is usually performed under a pressure of from normal pressure to 200 bar, preferably from normal pressure to 30 bar.

[0092] The temperature of the telomerisation reaction is usually in the range of 20 to 200 °C preferably of 30 to 180 °C, more preferably of 40 to 140 °C and still more preferably of 40 to 120 °C.

[0093] The concentration of the transition metal is usually in the range of 1 to 200 mM per litre of ionic liquid and more preferably 10 to 50.

[0094] The amount of the salt in the reaction system is usually 5 to 100 parts per 100 parts of the diolefin, preferably 10 to 50.

[0095] The molar ratio of telogen/conjugated diolefin is usually in the range 0.5 to 10, preferably 0.5 to 5.

10 [0096] As generally known, the telomerisation of butadiene with water as the telogen is preferably carried out in the presence of a base and under CO₂-pressure. Suitable bases include for example the hydroxides of alkali metals and alkaline earth metals and amines. Suitable amines include for example tertiary amines having a basicity constant (pK_a) of at least 7, for example tri(C1-C6)alkylamines such as
15 trimethylamine, triethylamine, tri-n-propylamine, tri-n-butylamine, etc.; aminoalcohols such as 1-N,N-dimethylamino-2-propanol, 1-N,N-dimethylamino-3-butanol, etc.; and N,N-dimethyl-2-methoxyethylamine, N,N-dimethyl-3-ethoxypropylamine, N-methylpyrrolidine, N-methylpiperidine, N-methylmorpholine, N,N'-dimethylpiperazine, N,N,N',N'-tetramethyl-1,3-butanediamine and the like.

20 Among these, triethylamine is most preferred.

[0097] Further carbonate and/or bicarbonate ions may be present along with the tertiary amine to accelerate the rate of n-octadienol formation. Carbonate and bicarbonate ions are conveniently derived from carbon dioxide, sodium bicarbonate or formic acid, which releases these ions in the reaction system. Among these, carbon
25 dioxide is most preferred.

[0098] The amount of carbon dioxide which promotes the butadiene telomerisation is not critical and may range from about 10⁻³ to 1, preferably from 10⁻² to 0.5, mol per mole of the conjugated diene.

[0099] The substituted 1,2,3-, 1,2,3,4- and 1,2,3,4,5- substituted imidazolium salts
30 of the present invention can be used as a solvent for the hydrodimerisation with rates under the conditions already known.

[00100] In the following examples the use according to the present invention is exemplified in the catalytic conversion of conjugated dienes (telomerisation) and oligomerisation of dienes. It is however to be understood that the present invention is not limited to such uses.

5 Fig. 1 shows the apparatus for telomerising butadiene which has been used in the examples.

 Fig. 2 shows an embodiment of a two-phase process of telomerisation of butadiene (C_4H_6) with MeOH or ethylene glycol in the ionic liquids in accordance with the present invention.

10

Examples

A- Preparation of non-aqueous ionic liquids

a- Materials

15 All syntheses are carried out under dry argon using standard Schlenk techniques. Methylene dichloride is distilled over P_2O_5 and stored over 0.3 nm molecular sieves. All other reagents (1,2-dimethylimidazole, pyridine, HPF_6 , 1-chlorobutane) are purchased from Aldrich and used as is, unless otherwise indicated.

20

b- Physicochemical measurements

 Proton and carbon NMR are recorded on a Bruker AC 300 MHz using CD_2Cl_2 (from SDS) as solvent and $SiMe_4$ (from Aldrich) as internal standard. Melting point of solid salts is measured on a digital apparatus from Electrothermal.

25

c- Syntheses

 The following salts are prepared according to known procedures (see e.g. for $[EMI][Br]$, $[EMI][Tf_2N]$, $[EMI][CF_3SO_3]$, $[BMI][Tf_2N]$: Bonhôte, P.; Dias, A.-P.; Papageorgiou, N.; Kalyanasundaram, K.; Gratzel, M. *Inorg. Chem.*, (1996), **35**, 1168-1178; for $[BMI][BF_4]$: Holbrey, J. D.; Seddon, K. R., *J. Chem. Soc., Dalton Trans.*, (1999), 2133-2139; for $[BMI][PF_6]$: Suarez, P. A. Z.; Dullius, J. E. L.; Einloft, S.; de Souza, R. F.; Dupont, J., *Polyhedron*, (1996), **15**, 1217-1219).

30

EMI⁺Br⁻ (1-ethyl-3-methylimidazolium bromide),

EMI⁺Tf₂N⁻ (1-ethyl-3-methylimidazolium bistrifluoromethylsulphonylamide),

5

EMI⁺CF₃SO₃⁻ (1-ethyl-3-methylimidazolium trifluoromethylsulphonate),

BMI⁺Cl⁻ (1-n-butyl-3-methylimidazolium chloride),

10

BMI⁺PF₆⁻ (1-n-butyl-3-methylimidazolium hexafluorophosphate),

BMI⁺Tf₂N⁻ (1-n-butyl-3-methylimidazolium bistrifluoromethylsulphonylamide),

and

BMI⁺BF₄⁻ (1-n-butyl-3-methylimidazolium tetrafluoroborate).

15

Preparation example 1

1-butyl-2,3-dimethylimidazolium chloride [BMMI][Cl]:

Freshly distilled 1-chlorobutane (88 g, 0.96 mol) is added in one portion to a 500 ml thick walled glass reactor equipped with a magnetic stirrer containing 1,2-dimethylimidazole (65 g, 0.68 mol). The reactor is sealed and the solution is stirred for 16 h at 100 °C (note: the reaction pressure is in excess of 2 atm.). Reactor is degassed and the hot solution is transferred (under argon) in a round bottom flask containing acetonitrile (95 ml). The solution is added drop wise under vigorous stirring to toluene (500 ml). A precipitate forms and is filtered out, washed with toluene (3 X 100 ml) and dried overnight under vacuum. [BMMI][Cl] is obtained (89.73 g, 70 % yield) as a white hygroscopic solid.

¹H NMR (CD₂Cl₂): δ 0.97 [t, ³J=7.15 Hz, NCH₂CH₂CH₂CH₃]; 1.39 [sext, ³J=7.5 Hz, NCH₂CH₂CH₂CH₃]; 1.80 [quint, ³J=7.4 Hz, NCH₂CH₂CH₂CH₃]; 2.75 [s, CCH₃]; 4.00 [s, NCH₃]; 4.19 [t, ³J=7.1 Hz, NCH₂CH₂CH₂CH₃]; 7.52 [s, CH]; 7.84 [s, CH];

¹³C NMR (CD₂Cl₂): δ 10.70 [NCH₂CH₂CH₂CH₃]; 13.74 [NCH₂CH₂CH₂CH₃]; 19.96

[NCH₂CH₂CH₂CH₃]; 32.26 [CCH₃]; 36.09 [NCH₃]; 48.94 [NCH₂CH₂CH₂CH₃];
121.72 [CH]; 123.64 [CH]; 144.03 [CCH₃];

Elemental analysis (calculated): %C=57.14 (57.29), %H=9.08 (9.08), %N=14.85 (14.85).

5 Mp = 104-105 °C.

Preparation example 2

1-butyl-2,3-dimethylimidazolium bistrifluoromethylsulphonylamide [BMMI][Tf₂N]:

A solution of lithium bis((trifluoromethyl)sulphonyl)amide (41.9 g, 0.149
10 mol) in 100 ml of H₂O is added drop wise to a solution of [BMMI][Cl] (24.32 g,
0.129 mol) in 150 ml of H₂O. The solution is stirred at 70 °C for 2 h and then cooled
to room temperature. Methylene dichloride (50 ml) is added, and all is transferred to
a separatory funnel. The lower phase (ionic liquid + CH₂Cl₂) is collected. Ionic liquid
is purified through a short alumina column, and the CH₂Cl₂ removed on a Rotavapor.
15 The resultant hydrophobic liquid is washed 3 times with 150 ml of H₂O and dried for
3 h at 50°C under vacuum to afford [BMMI][Tf₂N] (44.1 g, 78.9 % yield) as a
colourless liquid.

¹H NMR (CD₂Cl₂): δ 0.97 [t, ³J=7.15 Hz, NCH₂CH₂CH₂CH₃]; 1.37 [sext, ³J=7.5 Hz,
NCH₂CH₂CH₂CH₃]; 1.78 [quint, ³J=7.4 Hz, NCH₂CH₂CH₂CH₃]; 2.59 [s, CCH₃];
20 3.79 [s, NCH₃]; 4.04 [t, ³J=7.1 Hz, NCH₂CH₂CH₂CH₃]; 7.19 [s, CH]; 7.21 [s, CH];
¹³C NMR (CD₂Cl₂): δ 9.78 [NCH₂CH₂CH₂CH₃]; 13.53 [NCH₂CH₂CH₂CH₃]; 19.86
[NCH₂CH₂CH₂CH₃]; 31.90 [CCH₃]; 35.60 [NCH₃]; 49.00 [NCH₂CH₂CH₂CH₃];
121.37 [CH]; 122.91 [CH]; 144.21 [CCH₃]; 120.50 [q, J_{C-F} = 321.4 Hz,
(CF₃SO₂)₂N];

25 Elemental analysis (calculated): %C=30.68 (30.49), %H=3.96 (3.95), %N=9.66
(9.70), %Cl < 250 ppm (0)

Preparation example 3

1-butyl-2,3-dimethylimidazolium tetrafluoroborate [BMMI][BF₄]:

30 Procedure previously described for the synthesis of [BMMI][Tf₂N] is used
(except the washes with water). From 22.23 g (0.118 mol) of [BMMI][Cl] and

15.52 g (0.141 mol) of sodium tetrafluoroborate, there are obtained 25.33 g (90 % yield) of [BMMI][BF₄] as a colourless, very viscous and hydrophilic liquid.

¹H NMR (CD₂Cl₂): δ 0.96 [t, ³J=7.15 Hz, NCH₂CH₂CH₂CH₃]; 1.38 [sext, ³J=7.5 Hz, NCH₂CH₂CH₂CH₃]; 1.78 [quintet, ³J=7.4 Hz, NCH₂CH₂CH₂CH₃]; 2.60 [s, CCH₃];
5 3.80 [s, NCH₃]; 4.06 [t, ³J=7.1 Hz, NCH₂CH₂CH₂CH₃]; 7.25 [dd, CH]; 7.30 [dd, CH]; ¹³C NMR (CD₂Cl₂): δ 9.58 [NCH₂CH₂CH₂CH₃]; 13.61 [NCH₂CH₂CH₂CH₃];
19.79 [NCH₂CH₂CH₂CH₃]; 31.94 [CCH₃]; 35.36 [NCH₃]; 48.70 [NCH₂CH₂CH₂CH₃]; 121.30 [CH]; 122.88 [CH]; 144.31 [CCH₃].

Elemental analysis (calculated): %C=45.09 (45.03), %H=7.25 (7.14), %N=11.52
10 (11.67), %Cl=0.11 (0)

B- Catalysis

a- Materials

Pd(OAc)₂ (98 %) is purchased from Strem Chemicals, stored under argon and
15 used without further purification. Triphenylphosphine (PPh₃) is obtained from Aldrich, trisodium tris((*meta*-sulphonatophenyl)phosphine) (TPPTS) from the lab reserves, bis(p-sulphonatophenyl)phenylphosphine dihydrate dipotassium salt (TPPDS) from Strem Chemical and diphenylphosphinobenzene-3-sulphonic acid sodium salt (TPPMS) from TCI. Those phosphines are stored under argon and used
20 as such. Triethylamine (Fluka >98%) and dimethyldodecylamine (Aldrich 97%) are distilled under argon, freeze-pumped and degassed with argon just before use. Water used in all experiments is deionised and degassed by freeze-pumping and bubbling of argon. Methanol is obtained by refluxing over Mg/I₂ and stored over 3 Å molecular sieves. Anhydrous ethylene glycol is obtained from Aldrich. Heptane was distilled
25 over Na/K and stored over 3 Å molecular sieves under argon. Carbon dioxide (N45) and butadiene (N25) were obtained from "Air-Liquide" and used directly from cylinders.

b- Catalytic experiments and analyses: Catalytic runs:

30 All catalytic reactions are performed in a 100 ml magnetically stirred stainless steel autoclave equipped with an inner glass sleeve and an internal thermocouple

under argon atmosphere as shown in Figure 1. In a typical reaction, $\text{Pd}(\text{OAc})_2$, phosphine, ionic liquid, telogen (methanol, ethylene glycol or water) and eventually heptane and/or amine are introduced in a Schlenk and then transferred into the purged autoclave via a ball valve. The autoclave is cooled to $T \ll -10^\circ\text{C}$ and the
5 desired mass of butadiene is transferred from a lecture bottle resting on a scale (for telomerisation of butadiene with water reactions, this step is followed by the warming to room temperature of the autoclave in order to introduce 5 bar of carbon dioxide). The reactor is then heated to the desired reaction temperature. After the selected reaction time, the autoclave is cooled to 40°C and butadiene is condensed in
10 a volumetric glass cylinder cooled to -78°C . The volume of the liquid condensed in the cylinder (determined to be almost pure butadiene) is noted. The autoclave is then warmed to room temperature. The remaining liquid phases in the autoclave are recovered, weighed and further analyzed (GC, GC-MS and NMR).

Thus are obtained two measures of butadiene conversion, from the volume of
15 unreacted butadiene condensed in the graduated cylinder (Conv. A) and the mass increase of the liquid solution in the autoclave (Conv. B). Conversion A is very often higher than conversion B due to possible loss of butadiene during trapping. Exact conversion will be considered as the average between conversion A and conversion B.

20

Work-up:

When at the end of the run only a single liquid phase is present, a sample is analysed without further treatment. When multiple phases are present, only the upper phase is analysed for product.

25

Analyses:

The identification of the telomerisation products is carried out both by Hewlett-Packard GC/MS and NMR analyses. Finally they are quantitatively analysed by gas chromatography on a HP6890 chromatograph equipped with FID detector and
30 a HP1 column ($L = 30\text{ m}$, internal diameter = 0.32 mm , film thickness = $0.25\text{ }\mu\text{m}$). Injector temperature is 170°C and detector temperature is 180°C . The temperature

program is from 60°C (3 min.) to 100°C (0 min.) at a heating rate of 10°C/min. to 220°C (30 min.) at a heating rate of 5°C/min.

c- Data presentation

5 Abbreviations:

- OT : 1,3,7-octatriene and 1,3,6-octatriene
- VCH : 4-vinyl-cyclohexene
- Oligo. : products containing three and more units of butadiene
- 1-OMe : (*cis* + *trans*)-1-methoxy-2,7-octadiene
- 10 • 3-OMe : 3-methoxy-1,7-octadiene
- 1-Ol : 1-hydroxy-2,7-octadiene
- 3-Ol : 3-hydroxy-1,7-octadiene
- 1-OGly : (*cis* + *trans*) 1-hydroxy-2-(2,7-octadienyl-1-oxy) ethane
- 3-OGly : 1-hydroxy-2-(1,7-octadienyl-3-oxy) ethane
- 15 • (1+3)-Gly : bis(octadienyl-1-oxy)-1,2-ethane + bis(octadienyl-3-oxy)-1,2-ethane

Conversions and selectivities are defined by the following equations:

- 20 • Conv. A %: calculated from unreacted butadiene remaining after reaction

$$\text{Conv. A} = \frac{\text{mass of butadiene introduced} - \text{mass of butadiene recovered}}{\text{mass of butadiene introduced}}$$

- Conv. B %: calculated from the increase in the mass of the reactor

$$\text{Conv. B} = \frac{\text{increase of the mass of reactor contents}}{\text{mass of butadiene introduced}}$$

25

- GC selectivities toward products X are evaluated as:

$$\text{GC. Sel. (X)} = \frac{\text{moles of product X}}{\sum (\text{moles of products})}$$

- Selectivities to linear telomers (in the tables comprehensively reported as Ratio 1/3) are evaluated as:

$$\text{Ratio 1 / 3} = \frac{\text{moles of linear telomer}}{\text{moles of branched telomer}}$$

- 5
- Turn over number in mol C₄H₆/mol Pd is evaluated as:

$$\text{TON} = \frac{\text{mole of butadiene converted (average of Conv. A and Conv. B)}}{\text{moles of palladium}}$$

- Determination of the palladium leaching in the organic phase:

$$\% \text{ Pd leaching} = \frac{\text{mass of Pd in the organic phase}}{\text{mass of initial Pd}}$$

10

The analytical procedure consists in evaporating the organic products and solubilising the residue in a mixture of HNO₃/HCl. Palladium contained in this mixture is then quantified by ICP/SM.

15

d- Catalysis examples (Cat. Ex.) and Comparative catalysis examples (Comp. Cat. Ex.)

Using the telomerisation apparatus described above catalysis examples and comparative catalysis are carried out under the conditions indicated in the following.

Catalysis examples 1 to 3

Catalysis examples 1 to 3 for telomerising 1,3-butadiene with methanol are run using Pd(OAc)₂/PPh₃ as the catalyst system. Table 1 shows the results.

Table 1: Telomerisation of 1,3-butadiene with methanol in the presence of Pd(OAc)₂/PPh₃^(a)

Cat. Ex.	Ionic Liquid	Conv		time (h)	GC Sel. (mol%)				Ratio 1 / 3	TON
		A	B		1-OMe + 3-OMe	VCH	OT	Olig.		
1	[BMMI][BF ₄]	100	92	1	81.8	0.2	15.1	2.9	8	2619
2	[BMMI][Tf ₂ N]	100	89	1	84.0	0.2	13.7	2.1	9	2537

(a)- Reaction conditions: Pd(OAc)₂ = 30 mg (0.134 mmol); PPh₃ = 105 mg (0.400 mmol); Ionic liquid (4 ml); MeOH = 15 ml (370 mmol); C₄H₆ = 20 g (370 mmol); T = 85°C;

Excellent results are found for the 1,2,3-trialkylimidazolium ionic liquids regarding conversions and reaction rates (Cat. Ex. 1 and 2). Quantitative conversions are achieved after 1 h for the two solvents studied [BMMI][Tf₂N] and [BMMI][BF₄]. Selectivity toward telomers is around 83% and, again, there is a notable effect on the regioselectivity of the telomerisation reaction, the ratio 1-OMe/3-OMe in excess of 8-9. It seems that the nature of the anion of the salts (between the hydrophobic Tf₂N⁻ salt and the hydrophilic BF₄⁻ salt) does not induce any notable difference in terms of activity and selectivity. Furthermore, at the end of these reactions, absolutely no palladium black is observed. A two-phase liquid-liquid system remains in the case of 1,2,3-trialkylimidazolium salts. The lower phase corresponds to the ionic liquid whereas the upper one is a mixture of products and unreacted methanol.

Catalysis examples 3 to 5

Catalysis examples 3 to 5 are run using Pd(OAc)₂ and water-soluble TPPMS as catalyst system. The results are shown in table 2.

Table 2: Telomerisation of 1,3-butadiene with methanol in the presence of $\text{Pd}(\text{OAc})_2/\text{TPPMS}^{(a)}$

Cat. Ex.	$\text{Pd}(\text{OAc})_2$ (mmol)	Temp. (°C)	time (h)	Conv		GC Sel. (mol%)				Ratio 1 / 3	TON
				A	B	1-OMe + 3-OMe	VCH	OT	Oligomers		
3	0.134	85	1	100	93	77.9	0.3	17.1	4.7	13.7	2769
4	0.028	85	5.75	85	80	77.7	0.8	20.9	0.6	16.0	11092
5	0.025	110	5.75	89	75	50.4	3.0	41.8	4.8	11.7	11953

(a)- Reaction conditions: TPPMS = 3 eq/Pd; $[\text{BMMI}][\text{Tf}_2\text{N}]$ (4 ml); MeOH = 15 ml (370 mmol);

5 C_4H_6 = 20 g (370 mmol);

The use of TPPMs leads to single phase system at 93% conversion (Cat. Ex. 3). Selectivity toward telomer is lower with TPPMS than with TPP, 77.9 against 84.0%. However with TPPMS the ratio 1-OMe/3-OMe is largely better, 13.7. The addition of 16%wt (10ml) of heptane to the solution obtained in Cat. Ex. 3 leads to clean separation of an ionic liquid phase from the product/MeOH/heptane phase.

Catalysis examples 6 to 8

15 Catalysis examples 6 to 8 are run in the presence of n-heptane using $\text{Pd}(\text{OAc})_2$ and different phosphines. The palladium leaching into the organic phase is determined by microanalysis of the two phases. The results are shown in table 3.

Table 3: Telomerisation of 1,3-butadiene with methanol in the presence of n-heptane using $\text{Pd}(\text{OAc})_2/$ and different phosphines^(a)

Cat. Ex.	P	Conv		GC Sel. (mol%)				Ratio 1 / 3	% Pd leaching	TON
		A	B	1-OMe + 3-OMe	VCH	OT	Oligo.			
6	TPP	82	71	53.3	-	40.6	6.1	14.6	14.2	2127
7	TPPMS	74	66	70.3	0.6	14.1	15.0	13.1	1.9	1816
8	TPPDS	56	51	58.6	1.2	17.5	22.8	15.0	1.8	1373

(a)- Reaction conditions: $\text{Pd}(\text{OAc})_2$ = 30 mg (0.134 mmol); Phosphine = 3 eq/Pd; $[\text{BMMI}][\text{Tf}_2\text{N}]$ (4 ml); MeOH = 7.5 ml (185 mmol); heptane = 10 ml; C_4H_6 = 20 g (370 mmol); t = 3 h; T = 85 °C;

Catalysis examples 9 to 13

The product phase of catalysis example 7 is decanted and the palladium contained in the ionic liquid phase is reintroduced to the reactor with fresh reactants and co-solvent. Microanalysis of the organic phase at the end of each catalysis example 10 to 14 indicates 1-4% leaching of palladium to the organic phase. Slight changes in activity are observed with each cycle. At first, activities are actually higher (catalysis examples 10 to 12) perhaps as the initiation reactions (conversion of Pd(OAc)₂ to the active species) reach completion. Activity slowly degrades in catalysis examples 12 and 13, in part perhaps due to the mechanical difficulty of quantitatively recuperating the ionic liquid phase (drops of the liquid can remain in the Schlenk tube used for decantation and/or syringe), palladium leaching, or to some palladium deactivation. In no case is there obvious evidence for the formation of palladium black. The results are shown in table 4.

15

Table 4: Telomerisation of 1,3-butadiene with methanol in the presence of n-heptane using recycled ionic liquid phase ^(a)

Cat. Ex.	Conv		GC Sel. (mol%)				Ratio 1 / 3	TON
	A	B	1-OMe + 3-OMe	VCH	OT	Oligo.		
9	74	66	70.3	0.6	14.1	15.0	13.1	1816
10	82	73	72.2	0.5	13.2	14.2	12.4	2001
11	89	81	72.4	0.4	15.9	11.3	15.4	2249
12	80	68	75.6	0.5	16.2	7.7	16.3	1907
13	70	63	78.8	0.9	17.3	3.1	18.6	1709

(a)- Reaction conditions: Pd(OAc)₂ = 30 mg (0.134 mmol); TPPMS = 146 mg (0.401 mmol); [BMMI][Tf₂N] (4 ml); MeOH = 7.5 ml (185 mmol); heptane = 10 ml; C₄H₆ = 20 g (370 mmol); T = 85 °C; t = 3 h.

20

Comparative Catalysis Examples 1 to 5

Comparative catalysis examples 1 to 5 are run using 1,3-dialkylimidazolium salts. The results are shown in table 5.

25

Table 5: Telomerisation of 1,3-butadiene with methanol using Pd(OAc)₂/PPh₃ as a catalyst system in the presence of 1,3-dialkylimidazolium salts as ionic liquid^(a)

Comp. Cat.	Ionic	Conv	Conv	GC Sel. (mol%)				Ratio	TON
Ex.	Liquid	A	B	1-OMe + 3-OMe	VCH	OT	Oligo.	1/3	
1	[BMI][Tf ₂ N] ^(b)	10	7	62.0	18.4	3.5	16.1	7.8	241
2	[EMI][Tf ₂ N] ^(b)	7	3	66.1	21.6	1.7	10.6	5.8	146
3	[EMI][Tf ₂ N] ^(c)	16	6	27.5	44.7	0.6	27.1	6.5	271
4	[BMI][BF ₄] ^(c)	15	-1	49.6	33.9	1.2	15.4	5.1	165
5	[EMI][PF ₆] ^(d)	13	2	38.0	49.8	8.2	4.0	7	184

5 (a)- Reaction conditions: Pd(OAc)₂ = 30 mg (0.134 mmol); PPh₃ = 105 mg (0.400 mmol); MeOH = 15 ml (370 mmol); C₄H₆ = 20 g (370 mmol); T = 85 °C; t = 22 h. (b)- 1 ml of ionic liquid ; (c)- 4 ml of ionic liquid ; (d)- Ionic liquids purchased from Aldrich (97 % pure, white solid, equivalent of 1ml of ionic liquid)

10 Conversions are extremely low. There is, however, no obvious formation of palladium black: systems, at the end of the reaction, appear as homogeneous quite colourless liquids. The "palladium-phosphine" catalyst supposed to be formed *in situ* certainly suffered of some modifications and so deactivation. The possible contamination of halide impurities from the ionic liquid is first evoked. Indeed, 15 synthesis of some ionic liquids is based on the use of chloride intermediates, potentially present in the remaining salts. However, this hypothesis can be totally excluded to explain this phenomenon. Indeed the halide contents of ionic liquids are always under the detection limit (50-250 ppm) which would be too little for total contamination of the catalyst. Moreover, this behaviour is observed with both 20 commercial salts (Comp. Cat. Ex. 5) and ionic liquids synthesised in the laboratory which exclude any questions concerning purity of salts used in those experiments.

Comparative Catalysis Examples 6 to 9

25 Comparative catalysis examples 6 to 9 are run using 1,3-dialkylimidazolium salts and different phosphine/palladium ratios in order to determine their influence on the conversion and selectivity. The results are shown in table 6.

Table 6: Telomerisation of 1,3-butadiene with methanol using $\text{Pd}(\text{OAc})_2/\text{PPh}_3$ with different Pd/phosphine ratios as a catalyst system in the presence of 1,3-dialkylimidazolium salts as ionic liquid ^(a)

Comp. Cat.	P / Pd	Conv	Conv	GC Sel. (mol%)				Ratio	TON
Ex.	ratio	A	B	1-OMe + 3-OMe	VCH	OT	Oligo.	1 / 3	
6	3	11	1	56.2	32.9	1.7	9.2	5.9	166
7	4	11	5	72.9	27.1	-	-	5.9	228
8	10	26	19	73.0	9.9	-	17.1	6.5	593
9	20	30	23	71.0	12.7	2.3	14.0	6.5	745

(a)- Reaction conditions: $\text{Pd}(\text{OAc})_2$ = 30 mg (0.134 mmol); $[\text{EMI}][\text{Tf}_2\text{N}]$ (1ml); MeOH = 15 ml (370 mmol); C_4H_6 = 20 g (370 mmol); T = 85 °C; t = 22 h.

Again, even with a large excess of PPh_3 performances of the catalytic system remain very low. With 20 equivalents of phosphine per palladium, conversion does not exceed 26 % (Comp. Cat. Ex. 9).

Catalysis examples 14 and 15

Catalysis examples 14 to 15 for telomerising 1,3-butadiene with glycol in the presence of n-heptane are run using $\text{Pd}(\text{OAc})_2/\text{TPPMS}$ as the catalyst system. Table 7 shows the results.

Table 7: Telomerisation of 1,3-butadiene with Glycol in the presence of n-heptane using $\text{Pd}(\text{OAc})_2/\text{TPPMS}$ as a catalyst system ^(a)

Cat.	Conv	Conv	GC Sel. (mol%)					Ratio	TON
Exp.	A	B	1-OGly + 3-OGly	VCH	OT	Oligo.	(1+3)-OGly ²	1/3	
14 ^(b)	96	87	67.6	5.2	4.2	1.7	21.3	24	2767
15 ^(c)	93	81	79.9	-	6.9	-	13.2	26	2482

(a)- Reaction conditions: $\text{Pd}(\text{OAc})_2$ = 30 mg (0.134 mmol); TPPMS = 0.146 g (0.401 mmol); Glycol = 10.5 ml (185 mmol); $[\text{BMIM}][\text{Tf}_2\text{N}]$ = 4 ml; C_4H_6 = 20 g (370 mmol); T = 85 °C; t = 45 min.;

(b)- heptane 10 ml

(c)- heptane 30 ml.

Catalysis examples 16 and 17

- Catalysis example 16 for telomerising 1,3-butadiene with water are run using
5 Pd(OAc)₂/TPPMS as the catalyst system. Table 8 shows the results.

Table 8: Telomerisation of 1,3-butadiene with water using Pd(OAc)₂/TPPMS as a catalyst system ^(a)

Ref.	Amine	Heptane (ml)	Conv		% GC. (mol)				Ratio (1/3)	TON
			A	B	Octadienols	VCH	OT	Oligo.		
121 ^(b)	NEt ₃ ^(d)	-	95	95	74.0	-	18.8	7.2	17	1164
133 ^(c)	NEt ₃ ^(d)	30	78	70	67.9	<0.1	32.1	<0.1	18	961

- 10 (a)- Reaction conditions: Pd(OAc)₂ = 0.050 g (0.223 mmol); TPPMS = 324 mg (0.892 mmol); [BMMI][BF₄] = 4 ml; C₄H₆ = 15 g (278 mmol); H₂O = 5 g (278 mmol); P_{CO2} = 5 bar; T = 85 °C; (b)- t = 3 h; (c)- t = 5 h; (d)- NEt₃ = 50 eq/Pd;

Catalysis examples 18 and 21

15

Catalysis example 21 for dimerisation - oligomerisation of 1,3-butadiene are run using Pd(OAc)₂ as the catalyst system. Table 9 shows the results.

Table 9: Oligomerisation of 1,3-butadiene using Pd(OAc)₂ as a catalyst system ^(a)

Cat.	Ionic Liquid	Conv		% GC. (mol)				TON
		A	B	VCH	OT	Butadiene trimers	Olig.	
18	[EMI][Tf ₂ N]	41	34	29.9	10.6	39.2	20.3	463
19	[BMI][BF ₄]	9	8	52.4	9.6	8.6	29.4	106
20	[BMI][PF ₆]	38	33	34.7	8.7	28.3	28.3	520
21	[BMMI][Tf ₂ N]	70	67	70.9	6.3	10.7	12.1	727

- 20 (a)- Reaction conditions: Pd(OAc)₂ = 0.050 g (0.223 mmol); Ionic liquids = 4 ml; C₄H₆ = 15 g (278 mmol); H₂O = 5 g (278 mmol); P_{CO2} = 5 bar; T = 85 °C; t = 22 h;


In terms of butadiene conversion, one will note that the use of BMMI+Tf₂N- considerably increases the conversion.

5 The invention will now be described in greater detail with the aid of embodiments given as non-limiting examples.

CLAIMS

- 5 1. Use of 1,2,3-substituted imidazolium salts, 1,2,3,4-substituted imidazolium salts or 1,2,3,4,5-substituted imidazolium salts, with the exception of 1,2,3,4,5-pentamethylimidazolium-, 1,2,3,5-tetramethyl-4*H*-imidazolium-, 1,2,3,4-tetramethyl-5*H*-imidazolium, and 1-trimethylsilyl-2,3,5-trimethyl-4*H*-imidazolium-salts with sulphonated or carboxylated triesters of phosphorous
10 acid as anion, as solvent.
2. A use according to claim 1 wherein the 1,2,3-substituted imidazolium salts, 1,2,3,4-substituted imidazolium salts or 1,2,3,4,5-substituted imidazolium salts are of the following formula (1):

15



(1)

wherein R¹ and R³ are the same or different and are each selected from the group consisting of a chiral or achiral alkyl group having up to 10, preferably
20 up to 6 carbon atoms, a chiral or achiral cycloalkyl group of 4 to 10 carbon atoms, a chiral or achiral aryl group of 6 to 10 carbon atoms, and a chiral or achiral tri(C1-C10)alkylsilyl-group,

R² is selected from the group consisting of a chiral or achiral alkyl group having up to 10, preferably up to 6 carbon atoms, a chiral or achiral cycloalkyl
25 group of 4 to 10 carbon atoms, a chiral or achiral aryl group of 6 to 10 carbon atoms and a halogen atom; more preferably R² is an alkyl group, a cycloalkyl group or an aryl group as defined above,

R⁴ and R⁵ are the same or different and are each selected from the group consisting of hydrogen, a chiral or achiral alkyl group having up to 10, preferably up to 6 carbon atoms, a chiral or achiral cycloalkyl group of 4 to 10 carbon atoms, a chiral or achiral aryl group of 6 to 10 carbon atoms, a chiral or achiral tri(C1-C10)alkylsilyl group, and a halogen atom; more preferably R⁴ and R⁵ are hydrogen, an alkyl group, a cycloalkyl group, an aryl group, or a tri(C1-C10)alkylsilyl group, still more preferably hydrogen,

and X⁻ represents an anion.


3. A use of imidazolium salts of formula (1) according to claim 2 wherein:
- R¹, R² and R³, identical or different, are each independently chosen from (C1-C6)alkyl, (C1-C6)haloalkyl and cycloalkyl;
 - R⁴ and R⁵, identical or different, are each independently chosen from hydrogen, (C1-C6)alkyl, (C1-C6)haloalkyl and cycloalkyl;
 - and X⁻ represents an anion.
4. A use of imidazolium salts of formula (1) according to any of claims 2 or 3 wherein:
- R¹, R² and R³, identical or different, are each independently chosen from methyl, ethyl, propyl, *iso*-propyl, butyl, *sec*-butyl, 1-pentyl, 2-pentyl, 3-pentyl, *iso*-pentyl, hexyl, *iso*-hexyl, cyclohexyl, trifluoromethyl, 1,1,1-trifluoroethyl, perfluoroethyl, perfluoropropyl, 1,1,1,2,2,3,3-heptafluorobutyl, and perfluorobutyl;
 - R⁴ and R⁵, identical or different, are each independently chosen from hydrogen, methyl, ethyl, propyl, *iso*-propyl, butyl, *sec*-butyl, 1-pentyl, 2-pentyl, 3-pentyl, *iso*-pentyl, hexyl, *iso*-hexyl, cyclohexyl, trifluoromethyl, 1,1,1-trifluoroethyl, perfluoroethyl, perfluoropropyl, 1,1,1,2,2,3,3-heptafluorobutyl, and perfluorobutyl;
 - and X⁻ represents an anion as hereinbefore described.

5. A use according to any of claims 1 to 4, wherein X^- is selected from anions of the group consisting of fluoride, chloride, bromide, iodide, hexafluorophosphate, hexafluoroantimonate, hexafluoroarsenate, fluorosulphonate, tetrafluoroborate, bisperfluoro (C1-C10)alkylsulphonyl amides, perfluoro(C1-C10)alkyl sulphonates, dichlorocuprate, tetrachloroborate, tetrachloroaluminate, heptachlorodialuminate ($Al_2Cl_7^-$), trichlorozincate, and anions of sulphonated or carboxylated (alkyl and/or aryl)-phosphines or phosphites.
- 10 6. A use according to claims 1 to 5, wherein X^- is selected from anions of the group consisting of fluoride, chloride, bromide, iodide, hexafluorophosphate, fluorosulphonate, tetrachloroborate, tetrafluoroborate, bis-perfluoro(C1-C10)alkylsulphonyl amides, perfluoro(C1-C10)alkyl sulphonates, and anions of sulphonated or carboxylated (alkyl, cycloalkyl and/or aryl)-phosphines or
- 15 -phosphites.
7. A use according to any of the claims 1 to 6, wherein 1,2,3-tri(C1-C10)alkyl-imidazolium salts are used as a solvent.
- 20 8. A use according to any of the claims 1 to 7, wherein 1-ethyl-2,3-dimethylimidazolium bromide, 1-ethyl-2,3-dimethylimidazolium bistrifluoromethylsulphonylamide, 1-ethyl-2,3-dimethylimidazolium trifluoromethylsulphonate, 1-n-butyl-2,3-dimethylimidazolium chloride, 1-n-butyl-2,3-dimethylimidazolium hexafluorophosphate, 1-n-butyl-2,3-
- 25 dimethylimidazolium bistrifluoromethylsulphonylamide, and 1-n-butyl-2,3-methylimidazolium tetrafluoroborate are used as a solvent.
9. A use according to any of the claims 1 to 7, wherein a 1-n-butyl-2,3-dimethylimidazolium salt is used as a solvent.

10. A use according to any of claims 1 to 7 or 9, wherein 1-n-butyl-2,3-dimethylimidazolium chloride, 1-n-butyl-2,3-dimethylimidazolium hexafluorophosphate, 1-n-butyl-2,3-dimethylimidazolium bistrifluoromethylsulphonylamide and/or 1-n-butyl-2,3-methylimidazolium tetrafluoroborate are
5 used as solvents.
11. A use according to any of the claims 1 to 10, wherein the solvent is a solvent in organic reactions.
- 10 12. A use according to any of claims 1 to 11, wherein the organic reaction is selected from the group consisting of the catalytic conversion of olefins, the condensation reaction, the hydrogenation reaction, the isomerisation reaction, the Suzuki cross-coupling reactions, the amination reaction, the partial
15 oxidation of alkanes, the kinetic resolution of racemic mixtures, the hydrogenation of imines, the hydrogenation of ketones, the transfer hydrogenation and the hydroxylation of aromatic organic compounds.
13. A use according to any of claims 1 to 12, wherein the reaction of catalytic conversion of olefins is selected from the telomerisation of conjugated dienes,
20 the dimerisation of olefins, the oligomerisation of olefins, the polymerization of olefins, the alkylation of olefins, the hydrogenation of olefins, the olefin metathesis, the hydroformylation of olefins, the ring-closing metathesis of olefins, the ring-opening metathesis polymerisation of olefins, the symmetric or asymmetric epoxidation of olefins (including heteroatom substituted olefins)
25 and the cyclopropanation of olefins,
14. A use of 1,2,3-substituted imidazolium salts, 1,2,3,4-substituted imidazolium salts or 1,2,3,4,5-substituted imidazolium salts, with the exception of 1,2,3,4,5-pentamethyl imidazolium-, 1,2,3,5-tetramethyl-4*H*-imidazolium-, 1,2,3,4-tetramethyl-5*H*-imidazolium, and 1-trimethylsilyl-2,3,5-trimethyl-4*H*-
30 imidazolium salts with sulphonated or carboxylated triesters of phosphorous

acid as anion, preferably according to any of the claims 1 to 13 as a catalyst or catalyst component.

- 15 15. A use of 1,2,3-substituted imidazolium salts, 1,2,3,4-substituted imidazolium salts or 1,2,3,4,5-substituted imidazolium salts, with the exception of 1,2,3,4,5-pentamethylimidazolium-, 1,2,3-5-tetramethyl-4*H*-imidazolium-, 1,2,3,4-tetramethyl-5*H*-imidazolium-, and 1-trimethylsilyl-2,3,5-trimethyl-4*H*-imidazolium-salts with a sulphonated or carboxylated triester of phosphorous acid as anion, preferably according to any of the claims 1 to 14 in a two-phase
10 catalytic reaction.
16. Use of 1,2,3-substituted imidazolium salts, 1,2,3,4-substituted imidazolium salts or 1,2,3,4,5-substituted imidazolium salts with a metal-containing compound, preferably of transition metal in an organic reaction selected from
15 the group consisting of the catalytic conversion of olefins, in particular selected from the telomerisation of conjugated dienes, the dimerisation of olefins, the oligomerisation of olefins, the polymerization of olefins, the alkylation of olefins, the hydrogenation of olefins, the olefin metathesis, the hydroformylation of olefins, the ring-closing metathesis of olefins, the ring-
20 opening metathesis polymerisation of olefins, the symmetric or asymmetric epoxidation of olefins (including heteroatom substituted olefins) and the cyclopropanation of olefins, the condensation reaction, the hydrogenation reaction, the isomerisation reaction, the Suzuki cross-coupling reactions, the amination reaction, the partial oxidation of alkanes, the kinetic resolution of
25 racemic mixtures, the hydrogenation of imines, the hydrogenation of ketones, the transfer hydrogenation and the hydroxylation of aromatic organic compounds.
17. Use according to claim 16, wherein the salts are used as a solvent.
30
18. Imidazolium salts of the following formula (I):



(1)

wherein R^1 and R^3 are the same or different and are each selected from the group consisting of a chiral or achiral alkyl group having up to 10, preferably up to 6 carbon atoms, a chiral or achiral cycloalkyl group of 4 to 10 carbon atoms, a chiral or achiral aryl group of 6 to 10 carbon atoms, and a chiral or achiral tri(C1-C10)alkylsilyl group,

R^2 is selected from the group consisting of a chiral or achiral alkyl group having up to 10, preferably up to 6 carbon atoms, a chiral or achiral cycloalkyl group of 4 to 10 carbon atoms, a chiral or achiral aryl group of 6 to 10 carbon atoms and a halogen atom; more preferably R^2 is an alkyl group, a cycloalkyl group or an aryl group as defined above,

R^4 and R^5 are the same or different and are each selected from the group consisting of hydrogen, a chiral or achiral alkyl group having up to 10, preferably up to 6 carbon atoms, a chiral or achiral cycloalkyl group of 4 to 10 carbon atoms, a chiral or achiral aryl group of 6 to 10 carbon atoms, a chiral or achiral tri(C1-C10)alkylsilyl group, and a halogen atom; more preferably R^4 and R^5 are hydrogen, an alkyl group, a cycloalkyl group, an aryl group, or a tri(C1-C10)alkylsilyl group, still more preferably hydrogen,

and X^- represents an anion,

with the exception of 1,2,3,4,5-pentamethylimidazolium-, 1,2,3,5-tetramethyl-4*H*-imidazolium-, 1,2,3,4-tetramethyl-5*H*-imidazolium, and 1-trimethylsilyl-2,3,5-trimethyl-4*H*-imidazolium-salts of sulphonated or carboxylated triesters of phosphorous acid, 1,2,3,4,5-pentamethylimidazolium iodide, 1,2,3-trimethylimidazolium bromide, 1,2,3-trimethylimidazolium iodide, 1,2-dimethyl-3-ethylimidazolium bromide, 1,2-dimethyl-3-ethylimidazolium

chloride, 1,2-dimethyl-3-butylimidazolium fluoride, 1-butyl-2,3-dimethylimidazolium salt of 2,2'-(2,5-cyclohexadiene-1,4-ylidene)bis[propanedinitrile], 1,3-dimethyl-2,4,5-trichloro-imidazolium tetrafluoroborate, 1,3-dimethyl-2,4,5-tribromo-imidazolium tetrafluorofluoroborate, 1,3-dimethyl-2,4,5-tribromo-imidazolium bromide.

19. An imidazolium salt according to claim 18 wherein;

- R^1 , R^2 and R^3 , identical or different, are each independently chosen from (C1-C6)alkyl, (C1-C6)haloalkyl and cycloalkyl;

- R^4 and R^5 , identical or different, are each independently chosen from hydrogen, (C1-C6)alkyl, (C1-C6)haloalkyl and cycloalkyl;

- and X^- represents an anion.

with the exception of 1,2,3,4,5-pentamethylimidazolium-, 1,2,3,5-tetramethyl-4H-imidazolium-, and 1,2,3,4-tetramethyl-5H-imidazolium-salts of sulphonated or carboxylated triesters of phosphorous acid, 1,2,3,4,5-pentamethylimidazolium iodide, 1,2,3-trimethylimidazolium bromide, 1,2,3-trimethylimidazolium iodide, 1,2-dimethyl-3-ethylimidazolium bromide, 1,2-dimethyl-3-ethylimidazolium chloride, 1,2-dimethyl-3-butylimidazolium fluoride, 1-butyl-2,3-dimethylimidazolium salt of 2,2'-(2,5-cyclohexadiene-1,4-ylidene)bis[propanedinitrile], 1,3-dimethyl-2,4,5-trichloro-imidazolium tetrafluoroborate, 1,3-dimethyl-2,4,5-tribromo-imidazolium tetrafluorofluoroborate, 1,3-dimethyl-2,4,5-tribromo-imidazolium bromide.

20. An imidazolium salt according to any of claims 18 or 19, wherein:

- R^1 , R^2 and R^3 , identical or different, are each independently chosen from methyl, ethyl, propyl, *iso*-propyl, butyl, *sec*-butyl, 1-pentyl, 2-pentyl, 3-pentyl, *iso*-pentyl, hexyl, *iso*-hexyl, cyclohexyl, trifluoromethyl, 1,1,1-trifluoroethyl, perfluoroethyl, perfluoropropyl, 1,1,1,2,2,3,3-heptafluorobutyl, and perfluorobutyl;

- R^4 and R^5 , identical or different, are each independently chosen from hydrogen, methyl, ethyl, propyl, *iso*-propyl, butyl, *sec*-butyl, 1-pentyl, 2-pentyl,

3-pentyl, *iso*-pentyl, hexyl, *iso*-hexyl, cyclohexyl, trifluoromethyl, 1,1,1-trifluoroethyl, perfluoroethyl, perfluoropropyl, 1,1,1,2,2,3,3-heptafluorobutyl, and perfluorobutyl;

- and X⁻ represents an anion as hereinbefore described.

5 with the exception of 1,2,3,4,5-pentamethylimidazolium-, 1,2,3,5-tetramethyl-4*H*-imidazolium-, and 1,2,3,4-tetramethyl-5*H*-imidazolium-salts of sulphonated or carboxylated triesters of phosphorous acid, 1,2,3,4,5-pentamethylimidazolium iodide, 1,2,3-trimethylimidazolium bromide, 1,2,3-trimethylimidazolium iodide, 1,2-dimethyl-3-ethylimidazolium bromide, 1,2-
10 dimethyl-3-ethylimidazolium chloride, 1,2-dimethyl-3-butylimidazolium fluoride, 1-butyl-2,3-dimethylimidazolium salt of 2,2'-(2,5-cyclohexadiene-1,4-ylidene)bis[propanedinitrile], 1,3-dimethyl-2,4,5-trichloro-imidazolium tetrafluoroborate, 1,3-dimethyl-2,4,5-tribromo-imidazolium tetrafluorofluoroborate, 1,3-dimethyl-2,4,5-tribromo-imidazolium bromide.

15

21. An imidazolium salt according to any of claims 18 to 20, wherein X⁻ is selected from anions of the group consisting of fluoride, chloride, bromide, iodide, hexafluorophosphate, hexafluoroantimonate, hexafluoroarsenate, fluorosulphonate, tetrafluoroborate, bisperfluoro (C1-C10)alkylsulphonyl
20 amides, perfluoro(C1-C10)alkyl sulphonates, dichlorocuprate, tetrachloroborate, tetrachloroaluminate, heptachlorodialuminate (Al₂Cl₇⁻), trichlorozincate, and anions of sulphonated or carboxylated (alkyl and/or aryl)-phosphines or phosphites.

with the exception of 1,2,3,4,5-pentamethylimidazolium-, 1,2,3,5-tetramethyl-
25 4*H*-imidazolium-, 1,2,3,4-tetramethyl-5*H*-imidazolium, and 1-trimethylsilyl-2,3,5-trimethyl-4*H*-imidazolium-salts of sulphonated or carboxylated triesters of phosphorous acid, 1,2,3,4,5-pentamethylimidazolium iodide, 1,2,3-trimethylimidazolium bromide, 1,2,3-trimethylimidazolium iodide, 1,2-dimethyl-3-ethylimidazolium bromide, 1,2-dimethyl-3-ethylimidazolium
30 chloride, 1,2-dimethyl-3-butylimidazolium fluoride, 1-butyl-2,3-dimethylimidazolium salt of 2,2'-(2,5-cyclohexadiene-1,4-ylidene)bis[propane-

dinitrile], 1,3-dimethyl-2,4,5-trichloro-imidazolium tetrafluoroborate,
 1,3-dimethyl-2,4,5-tribromo-imidazolium tetrafluorofluoroborate,
 1,3-dimethyl-2,4,5-tribromo-imidazolium bromide.


- 5 22. An imidazolium salt according to any of claims 18 to 21 wherein X⁻ is selected from anions of the group consisting of fluoride, chloride, bromide, iodide, hexafluorophosphate, fluorosulphonate, tetrachloroborate, tetrafluoroborate, bis-perfluoro(C1-C10)alkylsulphonyl amides, perfluoro(C1-C10)alkyl
 10 sulphonates, and anions of sulphonated or carboxylated (alkyl, cycloalkyl and/or aryl)-phosphines or -phosphites.
 with the exception of 1,2,3,4,5-pentamethylimidazolium-, 1,2,3,5-tetramethyl-4*H*-imidazolium-, 1,2,3,4-tetramethyl-5*H*-imidazolium, and 1-trimethylsilyl-2,3,5-trimethyl-4*H*-imidazolium-salts of sulphonated or carboxylated triesters of phosphorous acid, 1,2,3,4,5-pentamethylimidazolium iodide, 1,2,3-
 15 trimethylimidazolium bromide, 1,2,3-trimethylimidazolium iodide, 1,2-dimethyl-3-ethylimidazolium bromide, 1,2-dimethyl-3-ethylimidazolium chloride, 1,2-dimethyl-3-butylimidazolium fluoride, 1,3-dimethyl-2,4,5-trichloro-imidazolium tetrafluoroborate, 1,3-dimethyl-2,4,5-tribromo-imidazolium tetrafluorofluoroborate, 1,3-dimethyl-2,4,5-tribromo-imidazolium
 20 bromide.
23. An imidazolium salt according to any of claims 18 to 22 selected from 1-ethyl-2,3-dimethylimidazolium bromide, 1-ethyl-2,3-dimethylimidazolium
 25 bistrifluoromethylsulphonylamide, 1-ethyl-2,3-dimethylimidazolium trifluoromethylsulphonate, 1-n-butyl-2,3-dimethylimidazolium chloride, 1-n-butyl-2,3-dimethylimidazolium hexafluorophosphate, 1-n-butyl-2,3-dimethylimidazolium bistrifluoromethylsulphonylamide, and 1-n-butyl-2,3-methylimidazolium tetrafluoroborate
- 30 24. An imidazolium salt according to any of claims 18 to 22, selected from 1-n-butyl-2,3-dimethylimidazolium bistrifluoro(C1-C10)alkylsulphonylamide, 1-n-


butyl-2,3-dimethylimidazolium tetrafluoroborate or 1-n-butyl-2,3-dimethylimidazolium hexafluorophosphate.

- 5 25. Composition comprising a 1,2,3- or 1,2,3,4- or 1,2,3,4,5- substituted imidazolium salt as solvent, a metal compound, and a phosphorous-containing compound.
- 10 26. Composition comprising a transition metal and a 1,2,3- or 1,2,3,4- or 1,2,3,4,5- substituted imidazoline salt, with the exception of 1,2,3,4,5- pentamethylimidazolium-, 1,2,3,5-tetramethyl-4*H*-imidazolium-, 1,2,3,4-tetramethyl-5*H*-imidazolium, and 1-trimethylsilyl-2,3,5-trimethyl-4*H*-imidazolium-salts with sulphonated or carboxylated triesters of phosphorous acid as anion.

1/2

Figure 1





INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 01/02003

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D233/54 C07F5/02 C07C307/02 C08F4/00 C08F36/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07F C07C C08F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 2 150 740 A (ALLIED CORPORATION) 3 July 1985 (1985-07-03) example 3	18
X	FR 2 434 156 A (FIRMA TH. GOLDSCHMIDT A.G.) 21 March 1980 (1980-03-21) claim 1	18
X	FR 2 380 732 A (BAYER AG) 15 September 1978 (1978-09-15) * active substance nr 4,11,19 and 30 *	18
X	FR 2 302 301 A (FIRMA TH. GOLDSCHMIDT A.G.) 24 September 1976 (1976-09-24) example 11	18

-/-

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

1 March 2002

Date of mailing of the international search report

08/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/IB 01/02003

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FR 2 303 802 A (FIRMA TH. GOLDSCHMIDT A.G.) 8 October 1976 (1976-10-08) examples 4-6	18
X	----- CHEMICAL ABSTRACTS, vol. 70, no. 3, 20 January 1969 (1969-01-20) Columbus, Ohio, US; abstract no. 11699f, SAWA, NATSUO ET AL.: "Quaternary imidazolium compounds." XP002169536 abstract -& DATABASE CHEMICAL ABSTRACTS 'Online! CA 70:11699, XP002169538 compounds with RN 21054-86-4, -78-4, -79-5 and -82-0 & JP 06 812354 A (TOHO RAYON CO., LTD.)	18
X	----- CHEMICAL ABSTRACTS, vol. 130, no. 15, 12 April 1999 (1999-04-12) Columbus, Ohio, US; abstract no. 197145h, GOTO, KENICHI ET AL.: "Lactic acid and its oligomers, high-molecular-weight poly(lactic acid), and high-efficiency manufacture." XP002169537 abstract -& DATABASE CHEMICAL ABSTRACT 'Online! CA 130:197145, XP002169539 compound with RN 125376-11-6 and its use as a catalyst & JP 01 135662 A (MITSUI CHEMICALS INC.)	14,18
X	----- US 5 827 602 A (VICTOR R. KOCH ET AL.) 27 October 1998 (1998-10-27) examples 1,5	18
X	----- WO 99 40025 A (ACEP INC.) 12 August 1999 (1999-08-12) example 3	18
X	----- US 5 683 832 A (PIERRE BONHOTE ET AL.) 4 November 1997 (1997-11-04) column 4, line 7 - line 8	18
A	----- WO 95 21871 A (BP CHEMICALS LTD) 17 August 1995 (1995-08-17) page 1, line 1 - line 5; claims	1,14

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 18-24(partly)

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty of compounds of general formula (I) of claim 18 (imidazolium salts "per se"). So many documents were retrieved that it is impossible to determine which parts of the claim 18 may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claim 18 is impossible. However with respect to the preparation examples of the present application the search can be considered complete.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 01/02003

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 2150740	A	03-07-1985	DE 3442172 A1 JP 61133580 A	05-06-1985 20-06-1986
FR 2434156	A	21-03-1980	DE 2829137 A1 AT 361137 B AT 720978 A AU 519753 B2 BE 872502 A4 CH 637942 A5 ES 476626 A2 FR 2434156 A2 GB 2024805 A ,B IT 1107982 B JP 1406665 C JP 55022652 A JP 62012201 B NL 7810275 A ,B, SE 443976 B SE 7812918 A US 4189311 A	17-01-1980 25-02-1981 15-07-1980 17-12-1981 30-03-1979 31-08-1983 01-07-1979 21-03-1980 16-01-1980 02-12-1985 27-10-1987 18-02-1980 17-03-1987 07-01-1980 17-03-1986 04-01-1980 19-02-1980
FR 2380732	A	15-09-1978	DE 2706839 A1 AU 3328778 A BE 864012 A1 BR 7800928 A DK 68878 A FR 2380732 A1 GB 1569240 A JP 53101534 A NL 7801755 A US 4150967 A ZA 7800928 A	24-08-1978 23-08-1979 16-08-1978 02-01-1979 18-08-1978 15-09-1978 11-06-1980 05-09-1978 21-08-1978 24-04-1979 31-01-1979
FR 2302301	A	24-09-1976	DE 2508420 A1 BE 838451 A1 CH 588811 A5 FR 2302301 A1 NL 7601609 A	09-09-1976 28-05-1976 15-06-1977 24-09-1976 31-08-1976
FR 2303802	A	08-10-1976	DE 2510525 A1 AT 350550 B AT 176776 A AT 353984 B AT 386478 A AU 1143276 A BE 839038 A1 CH 618688 A5 ES 445928 A1 FR 2303802 A1 GB 1503077 A IT 1057933 B JP 51115476 A NL 7601827 A ,B, SE 421419 B SE 7602144 A US 4062965 A	16-09-1976 11-06-1979 15-11-1978 10-12-1979 15-05-1979 01-09-1977 16-06-1976 15-08-1980 01-05-1977 08-10-1976 08-03-1978 30-03-1982 12-10-1976 14-09-1976 21-12-1981 13-09-1976 13-12-1977
US 5827602	A	27-10-1998	EP 0839139 A1	06-05-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Patent Application No

PCT/IB 01/02003

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5827602	A		JP 2001517205 T WO 9702252 A1	02-10-2001 23-01-1997
WO 9940025	A	12-08-1999	WO 9940025 A1 EP 0968181 A1 EP 0971854 A1 WO 9928292 A1 JP 2002500678 T JP 2001527505 T US 2002009635 A1	12-08-1999 05-01-2000 19-01-2000 10-06-1999 08-01-2002 25-12-2001 24-01-2002
US 5683832	A	04-11-1997	CH 688236 A5 FR 2728569 A1 CN 1132746 A ,B EP 0718288 A1 JP 8259543 A	30-06-1997 28-06-1996 09-10-1996 26-06-1996 08-10-1996
WO 9521871	A	17-08-1995	AU 1584895 A BR 9505775 A CA 2159479 A1 CN 1123031 A CZ 9502576 A3 EP 0693088 A1 FI 954807 A WO 9521871 A1 JP 8509242 T NO 954015 A ZA 9501060 A	29-08-1995 27-02-1996 17-08-1995 22-05-1996 17-01-1996 24-01-1996 09-10-1995 17-08-1995 01-10-1996 09-10-1995 12-08-1996

THIS PAGE BLANK (USPTO)